



# The stable isotopic signature of biologically produced molecular hydrogen (H<sub>2</sub>)

S. Walter<sup>1</sup>, S. Laukenmann<sup>2</sup>, A. J. M. Stams<sup>3</sup>, M. K. Vollmer<sup>4</sup>, G. Gleixner<sup>5</sup>, and T. Röckmann<sup>1</sup>

<sup>1</sup>Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University, Utrecht, The Netherlands

<sup>2</sup>Department of Atmospheric Chemistry, MPI for Chemistry, Mainz, Germany

<sup>3</sup>Laboratory of Microbiology, Wageningen University, Wageningen, The Netherlands

<sup>4</sup>Empa, Swiss Federal Laboratories for Materials Science and Technology, Laboratory for Air Pollution and Environmental Technology, Dübendorf, Switzerland

<sup>5</sup>Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena, Germany

Correspondence to: S. Walter (s.walter@uu.nl)

Received: 23 November 2011 – Published in Biogeosciences Discuss.: 22 December 2011

Revised: 16 May 2012 – Accepted: 29 September 2012 – Published: 24 October 2012

**Abstract.** Biologically produced molecular hydrogen (H<sub>2</sub>) is characterised by a very strong depletion in deuterium. Although the biological source to the atmosphere is small compared to photochemical or combustion sources, it makes an important contribution to the global isotope budget of H<sub>2</sub>. Large uncertainties exist in the quantification of the individual production and degradation processes that contribute to the atmospheric budget, and isotope measurements are a tool to distinguish the contributions from the different sources. Measurements of  $\delta D$  from the various H<sub>2</sub> sources are scarce and for biologically produced H<sub>2</sub> only very few measurements exist.

Here the first systematic study of the isotopic composition of biologically produced H<sub>2</sub> is presented. In a first set of experiments, we investigated  $\delta D$  of H<sub>2</sub> produced in a biogas plant, covering different treatments of biogas production. In a second set of experiments, we investigated pure cultures of several H<sub>2</sub> producing microorganisms such as bacteria or green algae. A Keeling plot analysis provides a robust overall source signature of  $\delta D = -712\text{‰} (\pm 13\text{‰})$  for the samples from the biogas reactor (at 38 °C,  $\delta D_{\text{H}_2\text{O}} = +73.4\text{‰}$ ), with a fractionation constant  $\varepsilon_{\text{H}_2\text{-H}_2\text{O}}$  of  $-689\text{‰} (\pm 20\text{‰})$  between H<sub>2</sub> and the water. The five experiments using pure culture samples from different microorganisms give a mean source signature of  $\delta D = -728\text{‰} (\pm 28\text{‰})$ , and a fractionation constant  $\varepsilon_{\text{H}_2\text{-H}_2\text{O}}$  of  $-711\text{‰} (\pm 34\text{‰})$  between H<sub>2</sub> and the water. The results confirm the massive deuterium depletion of biologically produced H<sub>2</sub> as was predicted by the cal-

ulation of the thermodynamic fractionation factors for hydrogen exchange between H<sub>2</sub> and water vapour. Systematic errors in the isotope scale are difficult to assess in the absence of international standards for  $\delta D$  of H<sub>2</sub>.

As expected for a thermodynamic equilibrium, the fractionation factor is temperature dependent, but largely independent of the substrates used and the H<sub>2</sub> production conditions. The equilibrium fractionation coefficient is positively correlated with temperature and we measured a rate of change of 2.3 ‰/°C between 45 °C and 60 °C, which is in general agreement with the theoretical prediction of 1.4 ‰/°C.

Our best experimental estimate for  $\varepsilon_{\text{H}_2\text{-H}_2\text{O}}$  at a temperature of 20 °C is  $-731\text{‰} (\pm 20\text{‰})$  for biologically produced H<sub>2</sub>. This value is close to the predicted value of  $-722\text{‰}$ , and we suggest using these values in future global H<sub>2</sub> isotope budget calculations and models with adjusting to regional temperatures for calculating  $\delta D$  values.

## 1 Introduction

Molecular hydrogen (H<sub>2</sub>) is the second most abundant reduced compound in the atmosphere, after methane (CH<sub>4</sub>). With a global average mixing ratio of  $\sim 530$  ppb (parts-per-billion, nmole/mole) and an atmospheric lifetime of  $\sim 2$  yr, it is responsible for a large fraction of the chemical turnover of hydrogen atoms in the atmosphere and

contributes significantly to atmospheric chemistry (Novelli et al., 1999; Hauglustaine and Ehhalt, 2002; Rahn et al., 2003; Ehhalt and Rohrer, 2009). By reaction with the hydroxyl radical ( $\bullet\text{OH}$ ), hydrogen indirectly increases the atmospheric lifetimes of other trace gases that also react with  $\bullet\text{OH}$ , for example, CH<sub>4</sub> and carbon monoxide (CO) (Prather, 2003; Schultz et al., 2003; Jacobson et al., 2005; Jacobson, 2008) and, therefore, acts as an 'indirect' greenhouse gas. In the stratosphere, oxidation of H<sub>2</sub> is a source of water vapour, which is important for the radiative properties of the stratosphere and also forms the substrate for polar stratospheric clouds, which are key ingredients in the formation of the polar ozone holes (Tromp et al., 2003; Warwick et al., 2004; Feck et al., 2008; Jacobson, 2008).

H<sub>2</sub> is considered as a promising future energy carrier. It can be produced chemically, physically and biologically. The shortage, increase in cost and climate impact of fossil fuels leads to increased interest in sustainable and clean production of H<sub>2</sub>. One possible source to accommodate the expected energy demand might be biologically produced H<sub>2</sub>, e.g., via fermentation or photosynthesis.

Numerous studies in the past have addressed the global atmospheric budget of H<sub>2</sub>, but still none of the individual source or sink strengths is constrained to better than  $\pm 25\%$  (Ehhalt and Rohrer, 2009). Additional information is expected to come from the analysis of the H<sub>2</sub> isotopic composition ( $\delta D$ ), because the different sources of H<sub>2</sub> have a very different deuterium content.  $\delta D$  is defined as the relative deviation of the D/H ratio in a sample from the same ratio in the international reference material Vienna Standard Mean Ocean Water (VSMOW). Also the kinetic fractionation in the two main removal processes, soil deposition and reaction with OH, is different.

Tropospheric H<sub>2</sub> is enriched in deuterium with  $\delta D \sim +130\%$ , (Gerst and Quay, 2001; Rhee et al., 2006; Rice et al., 2010; Batenburg et al., 2011) compared to surface emissions from fossil fuel combustion and biomass burning ( $\delta D$  approximately  $-200\%$  and  $-300\%$ , respectively) (Gerst and Quay, 2001; Rahn et al., 2002; Röckmann et al., 2010a; Vollmer et al., 2010). As originally proposed by Gerst and Quay (2001) from budget closure, the photochemical sources of H<sub>2</sub> are also enriched in deuterium with  $\delta D$  between  $\sim +100\%$  and  $+200\%$ , (Rahn et al., 2003; Röckmann et al., 2003, 2010b; Feilberg et al., 2007; Nilsson et al., 2007, 2010; Pieterse et al., 2009). Biologically produced H<sub>2</sub> has the most exceptional isotopic composition. Biochemical reactions take place in the aqueous phase and, therefore, the isotopic composition of biologically produced H<sub>2</sub> should reflect the thermodynamic isotope equilibrium between H<sub>2</sub> and H<sub>2</sub>O. Bottinga (1969) calculated fractionation factors for isotope equilibrium in the system H<sub>2</sub> – water vapour. He predicts  $\epsilon_{\text{H}_2\text{-H}_2\text{O}}$  values for biologically produced H<sub>2</sub> of  $-737\%$  to  $-693\%$ , relative to the water, in the main biological relevant temperature range between 10 °C and 40 °C. Up to now only few individual measurements have been carried out to

experimentally determine the isotopic composition of biologically produced H<sub>2</sub> and confirm the extremely depleted values calculated by Bottinga (1969). Rahn et al. (2002) measured headspace samples from a jar of termites with a value of  $\delta D = -778\%$  at a mixing ratio of 1.8 ppm (parts-per-million,  $\mu\text{mole/mole}$ ), and  $\delta D = -690\%$  at a mixing ratio of 4 ppm from a water headspace sample taken from an eutrophic pond. For none of these values the isotopic composition of the water was reported and it appears that the equilibrium isotope effect between H<sub>2</sub> and H<sub>2</sub>O has never been experimentally verified. Today recent global modelling studies have incorporated biological sources with an isotopic composition of  $\delta D = -628\%$  (Price et al. 2007; Pieterse et al. 2011).

Although biologically produced H<sub>2</sub> is only responsible for approximately 10% of the annual global H<sub>2</sub> source (Novelli et al., 1999; Hauglustaine and Ehhalt, 2002; Ehhalt and Rohrer, 2009; Pieterse et al., 2011) the extreme deuterium depletion relative to ambient atmospheric H<sub>2</sub> makes it a quite important contributor to the isotope budget (Price et al. 2007; Pieterse et al. 2011). An increasing demand and anthropogenic biological production of H<sub>2</sub> by e.g., industrial fermentation of biogenic waste material is associated with an expected release to the atmosphere because of leakage during production, storage, transport and use. This may increase the contribution of highly deuterium-depleted H<sub>2</sub> to the atmosphere.

Here we present the first systematic experimental evaluation of the isotope source signature of biologically produced H<sub>2</sub>, which is then compared to the theoretical calculations of Bottinga (1969). We measured the isotopic composition of fermentative produced molecular H<sub>2</sub> in biogas, using different production conditions and substrates. Additionally we investigated H<sub>2</sub> produced from pure cultures of fermentative bacteria (*Caldicellulosiruptur saccharolyticus*, *Escherichia coli*, and *Clostridium acetobutylicum*) and of one N<sub>2</sub>-fixer (*Azospirillum brasiliensis*). We also measured photosynthetically produced H<sub>2</sub> from the common green algae *Chlamydomonas reinhardtii*.

## 2 Experimental

### 2.1 Samples

#### 2.1.1 Samples from a biogas plant

Samples were provided from a biogas plant in Freising, Germany, where also the experiments were conducted. Experiments were carried out with batch cultures (2 l Merck glass bottles) and continuous cultures (30 l glass container). Both were fed with different substrates from surrounding agriculturalures such as corn, sunflower, cellulose, grass, wheat or mixtures of these substrates. For both treatments the same inoculum was used. It was provided from a pilot-plant scaled plant

(3500 l volume). An overview about used substrates and different treatments is given in Table 1.

The batch cultures consist of 1600 ml inoculum and were fed once with 50 g substrate (organic dry substance, oDS) and incubated at a stable temperature of 38 °C. After 35 days, headspace gas samples were taken with gas tight syringes into evacuated 12 ml glass tubes with an overpressure of approximately 1 bar.

The continuous cultures consist of 30 l inoculum and were fed daily in the morning and incubated at temperatures of 38 °C to 60 °C depending on the treatment. The treatments also differ in the amount of substrate between 2 and 3.5 kg organic dry substance/day (oDS/d). Approximately 4 h after feeding, samples were taken at a syringe port at the fermenter with gas tight syringes into evacuated 12 ml glass tubes with an overpressure of approximately 1 bar.

In total, three samples from batch cultures and 13 samples from continuous cultures were measured (see Table 1). Some samples were measured in duplicate or more. The headspace of pure inoculum was also sampled and measured.

### 2.1.2 Pure culture experiments

Headspace samples were taken from 5 pure cultures of known H<sub>2</sub> producing organisms: (i) three common fermentative bacteria: *Caldicellulosiruptur saccharolyticus*, *Escherichia coli* and *Clostridium acetobutylicum*; (ii) one N<sub>2</sub>-fixing bacterium (*Azospirillum brasiliensis*, strain SP7); and (iii) one limnic green algae *Chlamydomonas reinhardtii* (SAG strain number 11–32b).

*E. coli* and *C. saccharolyticus* were grown with 10 or 20 mM glucose and 0.2 g l<sup>-1</sup> yeast extract in the medium as described in Stams et al. (1993). These bacteria were grown in 120-ml vials with 50 ml medium or 250-ml bottles with 100 ml medium, and a gasphase of N<sub>2</sub>/CO<sub>2</sub> (80/20). *E. coli* was grown at 37 °C and *C. saccharolyticus* at 70 °C. *C. acetobutylicum* was grown at 37 °C as described by Nimcevic et al. (1998). The gas phase was N<sub>2</sub>. Gas samples were taken from the cultures at the end of growth by gastight syringes and injected in sterile vacuum vials, previously flushed with pure nitrogen.

12 ml of preincubated *A. brasiliensis* (strain SP7) was used to inoculate 600 ml of ampicillin medium in a closed 2 l borosilicate bottle. Three replicates and one control were incubated for 5 days at 30 °C. The headspace gas volume was sampled into a pre-evacuated 1 l glass container (NORMAG, Illmenau, Germany) sealed with two polychlorotrifluoroethylene (PCTFE) valves.

The green algae *C. reinhardtii* was cultivated in a sulfate-limited Tris-Acetate-Phosphate (TAP) medium as part of an experiment conducted in Switzerland and described in more detail by Haus et al. (2009). For the batch used in our isotope study, a N<sub>2</sub> headspace technique in a glass bottle was applied. After approximately 8 days of incubation, several ml of headspace gas were extracted using a gastight syringe

and injected into a pre-evacuated 1 l glass container of the same type as mentioned above. Synthetic air, further purified from traces of H<sub>2</sub> using a catalyst (Sofnocat 514, Molecular Products, Thaxted, UK) was added (to 1.9 bar total pressure) to dilute the sample and, thereby, making it suitable for H<sub>2</sub> measurements. Initial H<sub>2</sub> mixing ratio measurements were conducted at Empa before transferring the sample to IMAU for detailed H<sub>2</sub> and  $\delta D$  analysis. Results for H<sub>2</sub> mixing ratios of Empa and IMAU are in agreement within the error bars and the direct comparison is not shown here.

### 2.2 Determination of H<sub>2</sub> mixing ratio and isotopic composition

The mixing ratio and isotopic composition of molecular H<sub>2</sub> was determined by using the experimental setup developed by Rhee et al. (2004) and modified as described in Röckmann et al. (2010b). Due to a lack of international isotope standards for H<sub>2</sub>, calibration is a critical issue. Our calibration scale has been described in Batenburg et al. (2011). Samples were measured randomly and within 35 days after collection. The measurements consist of the following steps: (1) The sample is cryogenically separated at -240 °C, which means that all gaseous compounds, with the exception of H<sub>2</sub> and some noble gases, are condensed; (2) The non-condensed fraction of the sample (including H<sub>2</sub>) is preconcentrated using a 5 Å molecular sieve at -210 °C; (3) H<sub>2</sub> is focused on a capillary gas chromatographic column (5 Å molecular sieve) and chromatographically purified from remaining contaminants at 50 °C; (4) the D/H ratio of molecular H<sub>2</sub> is determined by continuous flow isotope ratio mass spectrometry using a ThermoFinnigan Delta Plus XL instrument.

The analytical system is designed for measurement of air samples with H<sub>2</sub> mixing ratios in the range of typical atmospheric air samples (e.g., Röckmann et al. 2003, 2010b; Rhee et al. 2006; Batenburg et al. 2011). The samples obtained from the biogas reactor and the individual cultures have extremely high H<sub>2</sub> molar mixing ratios between 10 ppm and 1.4 % (see Table 1), which are outside the measurement range of our instrument. To a certain degree, the analytical system has some flexibility as regards high H<sub>2</sub> mixing ratios because simply smaller samples can be inserted into the sample volume; however, in this study the values were that high that the samples had to be diluted. Two dilution methods were adapted for samples of pure cultures and biogas samples.

Several samples from the pure cultures were expanded into 2 l electropolished stainless steel canisters that are routinely used in our laboratory for airborne air sampling (Kaiser et al., 2006; Laube et al., 2008, 2010 and referenced herein). They were diluted by a factor of approximately 2000 with H<sub>2</sub>-free synthetic N<sub>2</sub>-O<sub>2</sub> mixtures. The mixtures were then measured as normal air samples. This procedure induces errors from the dilution itself (for the mixing ratios) and from unquantifiable blank levels of H<sub>2</sub> in the dilution gas. Another disadvantage is that no reference gases are available in the

**Table 1.** Molecular hydrogen (H<sub>2</sub>) mixing ratio and  $\delta D$  (vs. VSMOW) from different biogas production treatments and pure cultures. Columns 4 and 5 give the raw (i.e. measured) values for mixing ratio and  $\delta D$ , which are used in the Keeling plot in Fig. 1. Columns 6 and 7 give the final values of the pure sample after correction for the dilution with standard gas (with known H<sub>2</sub> mixing ratio of 546.2 ppb,  $\delta D + 71.4$  ‰, used for biogas samples) and H<sub>2</sub> free gas (used for pure microorganism culture samples), respectively. Substrate was added in units of kg organic dry material per day [o DM/d]. Some samples are measured in duplicate or more. Uncertainties in the table are given for the reproducibility of the measurements or calculations as the average of absolute deviation of data from their mean. For single measurements we assume a mean uncertainty of 4.1 % for mixing ratios and 2.4 % for  $\delta D$  values.

Culture	Temp. (°C)	Substrate (organic dry material/day)	Measured mixing ratio (ppb)	Measured $\delta D$ (‰)	Corrected mixing ratio (ppm)	Corrected $\delta D$ (‰)
inoculum	38	inoculum	624	-12	27	-587
batch culture	38	corn cob	584	+12	13	-831
	38	corn + sunflower	575	+35	10	-636
	38	cellulose	592	+21	16	-555
continuous culture	38	grass, 2 kg o DM/d	702	-104	53	-712
	38	grass, 2 kg o DM/d	1252	-370	240	-710
	38	grass, 2 kg o DM/d	1531	-437	335	-718
	38	corn, 2 kg o DM/d	675	-85	44	-741
	38	corn, 2.5 kg o DM/d	587	+16	14	-710
	38	corn, 3.5 kg o DM/d	689	-90	49	-699
	38	grass, 2 kg o DM/d ( $n = 2$ )	1170 ± 23	-350 ± 6	212 ± 5	-718 ± 21
	38	grass, 2 kg o DM/d ( $n = 3$ )	2262 ± 135	-520 ± 9	587 ± 38	-708 ± 20
	38	grass, 2 kg o DM/d	1133	-327	201	-696
	45	30 % grass + 30 % corn + 40 % cereals ( $n = 2$ )	946 ± 23	-270 ± 13	138 ± 7	-734 ± 22
	50	30 % grass + 30 % corn + 40 % cereals ( $n = 2$ )	1371 ± 87	-408 ± 15	282 ± 15	-726 ± 21
	55	30 % grass + 30 % corn + 40 % cereals ( $n = 2$ )	2273 ± 121	-523 ± 6	584 ± 32	-711 ± 21
	60	30 % grass + 30 % corn + 40 % cereals ( $n = 2$ )	2510 ± 66	-535 ± 7	671 ± 26	-703 ± 20
	Microorganism species					
Pure microorganism culture	70	<i>Caldicellulosiruptur saccharolyticus</i>	6360	-758	13887	-758
	37	<i>Escherichia coli</i> ( $n = 2$ )	3507 ± 9	-758 ± 0	8179 ± 21	-758 ± 0
	37	<i>Clostridium acetobutylicum</i>	11422	-741	13403	-741
	20	<i>Chlamydomonas reinhardtii</i>	596	-721	2285	-721
	30	<i>Azospirillum brasiliensis</i> ( $n = 5$ )	4043 ± 1096	-556 ± 24	1339 ± 420	-664 ± 19

region of the extremely deuterium-depleted samples, and the isotope scale has to be extrapolated very far outside the range that was used for calibrating the reference gas (-9.5 ‰ to +205 ‰) (Batenburg et al., 2011). Therefore, for the samples from the biogas plant and the N<sub>2</sub> fixer (*A. brasiliensis*), a standard addition method was developed. Small amounts of a sample (usually approximately 1 ml) were added manually with a gas tight syringe to air from the laboratory reference air cylinder (H<sub>2</sub> mixing ratio = 546.2 ppb,  $\delta D = +71.4$  ‰) (Batenburg et al., 2011). For the biogas samples following this procedure the measured mixing ratios after dilution were between 575 ppb and 2510 ppb, and  $\delta D$  values were between +35 ‰ and -535 ‰ (Table 1). This means that in the measurement procedure itself a “Keeling plot analysis” is involved, because the H<sub>2</sub> and HD measured in the isotope ratio mass spectrometer is then a mixture of the well-known reference air and the unknown sample (Fig. 1 for the biogas samples). The isotopic composition of the original sample is then inferred by extrapolation of the linear fit to the correlation

between  $\delta D$  and inverse mixing ratio to 0 (y-axis intercept). On the one hand, this introduces an error from the extrapolation, but on the other hand the measured  $\delta D$  values are much closer to the range that was used for calibration of the reference gas. The manual injection of the reference gas with a syringe leads to a relatively high error for the reproducibility of mixing ratios for the original biogas samples ( $\pm 4.1$  %), whereas the reproducibility for  $\delta D$  is not much worse than for normal atmospheric air samples ( $\pm 2.4$  %), since an error in mixing ratio only changes the location of the mixture on the mixing line, but not the y-axis intercept (see Fig. 1). The error for measurement reproducibility is given as the average of absolute deviation of data from their mean.

The isotopic composition of the water used in the incubation experiments was determined by Hydroisotop GmbH, Schweitenkirchen, Germany.

The  $\delta D$  is defined as followed:

$$\delta D = \frac{\left[ \frac{D}{H} \right]_{\text{sample}}}{\left[ \frac{D}{H} \right]_{\text{standard}}} - 1 \quad (1)$$

Fractionation constants  $\varepsilon_{\text{H}_2\text{-H}_2\text{O}}$  of H<sub>2</sub> relative to H<sub>2</sub>O were calculated as

$$\begin{aligned} \varepsilon_{\text{H}_2\text{-H}_2\text{O}} &= \alpha_{\text{H}_2\text{-H}_2\text{O}} - 1 = \frac{\left[ \frac{D}{H} \right]_{\text{H}_2}}{\left[ \frac{D}{H} \right]_{\text{H}_2\text{O}}} - 1 \\ &= \frac{\delta D_{\text{H}_2} + 1}{\delta D_{\text{H}_2\text{O}} + 1} - 1 = \frac{\delta D_{\text{H}_2} - \delta D_{\text{H}_2\text{O}}}{\delta D_{\text{H}_2\text{O}} + 1} \end{aligned} \quad (2)$$


where  $\alpha$  is the fractionation factor between the H<sub>2</sub> product and the H<sub>2</sub>O reactant. All  $\delta D$  values are reported relative to Vienna Standard Mean Ocean Water (VSMOW). H<sub>2</sub> mixing ratios are reported as molar mixing ratios in parts per million (ppm =  $\mu\text{mole/mole}$ ) or parts per billion (ppb =  $\text{nmole/mole}$ ), or in percent (%) for high mixing ratios.

### 3 Results and discussion

Table 1 provides a summary of the results. The H<sub>2</sub> content of the samples differed considerably. While the samples of the batch incubations from the biogas reactor contained relatively low H<sub>2</sub> mixing ratios with a maximum of 16 ppm (0.0016 %), the continuous incubations showed values up to 671 ppm (0.0671 %). In such biogas reactors, hydrogen produced by anaerobic bacteria is metabolised by the activity of methanogenic archaea, resulting in rather low hydrogen partial pressures (Stams and Plugge, 2009). The highest H<sub>2</sub> mixing ratios up to 13887 ppm or 1.4 % were measured in the headspace of the pure culture of *C. saccharolyticus*.

A Keeling plot analysis provides a powerful tool to establish a robust overall source signature for the samples from the biogas reactor. The results of all measurements can fit very well ( $R^2 = 0.999$ ) on a straight line with a y-axis intercept (source signature) of  $\delta D = -713 \text{‰}$  ( $\pm 13 \text{‰}$ ). It should be noted that this fit is not equally constrained by all samples, but more influenced by samples with high mixing ratios. To examine the effect of a possible nonlinearity in the isotope scale the analysis was repeated using only samples with  $\delta D$  values  $> -100 \text{‰}$ . This results in a source signature of  $-718 \text{‰}$ , which shows that nonlinearity of the isotope scale does not significantly affect our results.

Results for the individual samples are also included in Table 1. The samples from the batch incubations show individually a large degree of variability, but this is due to the low H<sub>2</sub> mixing ratios of the sample-reference gas mixtures measured (see Sect. 2). For individual samples, the Keeling plot analysis covers only a small range in inverse mixing ratio and the associated errors after extrapolation to 0 are large. The average of all individual source signature determinations is



**Fig. 1.** Keeling plot of the diluted samples from the biogas reactor at 38 °C (black dots, excluding the inoculum). The least-square fit result is  $y = 427.4x - 711.55$ ,  $R^2 = 0.999$ . For completeness the results of the treatments at different temperatures, of the inoculum and the pure cultures are also included in the figure (triangles), but not taken into account for determination of the source signature.

$\delta D = -706 \text{‰}$  ( $\pm 37 \text{‰}$ ,  $n = 22$ ). This is in very good agreement with the Keeling plot approach above, which provides the more robust constraint, because the influence of the samples with low mixing ratios is reduced in the Keeling plot approach.

The  $\delta D$  source signatures of the pure cultures range between  $-664 \text{‰}$  for *A. brasiliensis* to  $-758 \text{‰}$  for *E. coli* and *C. saccharolyticus* and give a mean source signature of  $\delta D = -728 \text{‰}$  ( $\pm 28 \text{‰}$ ) (Table 2). Overall, the results from the different experimental setups confirm the very depleted source signature of biological H<sub>2</sub>, but there are differences between the different experimental setups. For further interpretation, the  $\delta D$  values of the H<sub>2</sub> produced must be compared to the source water (Table 2). The isotopic composition of the waters used in the biogas plant in Freising was  $\delta D = -73.4 \text{‰}$  ( $\pm 1.5 \text{‰}$ ). For the cultivation of the microorganisms the isotopic composition of the used water was  $-65.6 \text{‰}$  ( $\pm 1.5 \text{‰}$ ) (*C. reinhardtii*, Switzerland) and  $-52.1 \text{‰}$  ( $\pm 1.5 \text{‰}$ ) (fermenters cultivated in Wageningen). The isotopic composition of the water used in the cultivation of *A. brasiliensis* in Munich was not determined, but is assumed to be close to typical tap water in Freising ( $\delta D = -73.4 \text{‰}$ ). With this information, the fractionation constant  $\varepsilon$  between H<sub>2</sub> and H<sub>2</sub>O can be calculated Eq. (2), which eliminates the water as free parameter. It is evident from the results in Table 2 that the variability between the different experimental setups remains when corrected for the different waters.

The fact that the H<sub>2</sub> samples from the pure cultures were measured after dilution, and the biogas samples with the on-line mixing approach is unlikely to cause the difference, as a potential contamination during dilution of the pure culture samples should not lead to even lower  $\delta D$  values for the mixtures. It was beyond the scope of this project to further

**Table 2.** Isotopic source signature of produced H<sub>2</sub> ( $\delta D_{H_2}$ ), isotopic composition of the water used in the incubation ( $\delta D_{H_2O}$ ), and isotopic fractionation constant  $\epsilon$  between H<sub>2</sub> and H<sub>2</sub>O,  $\epsilon_{H_2-H_2O}$ . The  $\delta D_{H_2}$  values for the biogas samples are taken from the Keeling plot, values of  $\epsilon$  from Bottinga (1969) are interpolated to the actual temperature. All values are in ‰.

Sample origin	$\delta D_{H_2}$	$\delta D_{H_2O}$	$\epsilon_{H_2-H_2O}$	Bottinga (1969) $\epsilon_{H_2-H_2O}$
Biogas (Keeling plot), 38 °C – 60 °C ( $n = 24$ )	$-713 \pm 13$	$-73.4 \pm 1.5$	$-691 \pm 20$	
Biogas (Keeling plot), 38 °C ( $n = 15$ )	$-712 \pm 13$	$-73.4 \pm 1.5$	$-689 \pm 20$	$-695$ ‰
<i>Azospirillum brasiliensis</i> , 30 °C ( $n = 5$ )	$-664 \pm 19$	$-73.4 \pm 1.5$	$-637 \pm 19$	$-707$ ‰
<i>Caldicellulosiruptur saccharolyticus</i> , 70 °C	$-758$	$-52.1 \pm 1.5$	$-745$	$-650$ ‰
<i>Escherichia coli</i> , 37 °C	$-758 \pm 0$	$-52.1 \pm 1.5$	$-745 \pm 3$	$-697$ ‰
<i>Clostridium acetobutylicum</i> , 37 °C ( $n = 2$ )	$-741$	$-52.1 \pm 1.5$	$-726$	$-697$ ‰
<i>Chlamydomonas reinhardtii</i> , 20 °C	$-721$	$-65.5 \pm 1.5$	$-701$	$-722$ ‰
Sample biogas (grass/corn/cereals) at incubation temperatures of				
45 °C ( $n = 2$ )	$-734 \pm 22$	$-73.4 \pm 1.5$	$-713 \pm 24$	$-685$
50 °C ( $n = 2$ )	$-726 \pm 21$	$-73.4 \pm 1.5$	$-704 \pm 24$	$-678$
55 °C ( $n = 2$ )	$-711 \pm 21$	$-73.4 \pm 1.5$	$-688 \pm 23$	$-671$
60 °C ( $n = 2$ )	$-703 \pm 20$	$-73.4 \pm 1.5$	$-680 \pm 21$	$-664$

investigate whether these differences are significant, but this would be an interesting task for the future. In the absence of further information, it may not be appropriate to simply average the results from this to some degree arbitrary selection of samples to obtain a representative mean.

The result from the biogas samples is best constrained, however, this value is determined for a temperature range of 38 °C to 60 °C. Including only biogas samples at 38 °C (inoculum and treatments at higher temperatures are excluded from the Keeling plot, Fig. 1) we end up with a  $\delta D$  of  $-712$  ‰  $\pm 13$  ‰ and a fractionation constant  $\epsilon_{H_2-H_2O}$  of  $-689$  ‰  $\pm 20$  ‰. This value is our best estimate for a fractionation constant  $\epsilon_{H_2-H_2O}$  at 38 °C. Although systematic errors in the isotopic scale cannot be excluded due to a lack of international isotope standards, the good agreement with the theoretically calculated value of  $\delta D = -695$  ‰ (Bottinga, 1969) provides strong support for the validity of our results.


Bottinga (1969) also reports the temperature dependence of  $\epsilon_{H_2-H_2O}$ . We determined this temperature dependence experimentally over the incubation range 45 °C–60 °C with otherwise identical conditions (same inoculum and substrate, 30 % grass, 30 % maize, 40 % cereals). As expected for an enzymatic-catalysed reaction in this temperature range, the mixing ratio of H<sub>2</sub> is increasing with increasing temperatures (Fig. 2a). Figure 2b shows that  $\epsilon_{H_2-H_2O}$  increases with increasing incubation temperature from  $-713$  ‰ at 45 °C to  $-680$  ‰ at 60 °C, thus, by 2.3 ‰/°C. Gray diamonds in Fig. 2b indicate the theoretically predicted temperature dependency from Bottinga (1969), which is slightly smaller with 1.4 ‰/°C over the same temperature range. The measurements show a distinct offset of 28 ‰ at 45 °C reducing to 16 ‰ at 60 °C, relative to the theoretical results over this temperature range (Fig. 2). This offset is slightly larger than our estimated experimental uncertainty and remains at present

unexplained. Possible contributing factors in the measurements are the potential errors in the absolute isotope calibration (Batenburg et al. 2011) or a nonlinearity in the isotope scale at very low  $\delta D$  values, which is not obvious from the Keeling plot. Nevertheless, the overall temperature dependence is in good qualitative agreement with the calculations of Bottinga (1969), and we conclude that the experimental techniques are sufficiently advanced now to detect such small changes in the region of very depleted isotope values.

In order to derive a revised  $\delta D$  value for H<sub>2</sub> from biological sources that can be used in global models or isotope budget calculations, we calculated  $\epsilon_{H_2-H_2O}$  at a mean temperature of 20 °C using the measured value at 38 °C and our experimentally determined temperature dependence, yielding a value of  $\epsilon_{H_2-H_2O} = -731$  ‰ ( $\pm 20$  ‰). For calculating a global average  $\delta D$  value of H<sub>2</sub> from biological sources we used a global average value of  $\delta D$  of precipitation of  $\delta D = -37.8$  ‰ (Hoffmann et al. 1998; Bowen and Revenaugh, 2003), and then calculate  $\delta D = -741$  ‰ ( $\pm 20$  ‰). Using the theoretical temperature dependence of Bottinga, we calculate  $\epsilon_{H_2-H_2O} = -715$  ‰ ( $\pm 20$  ‰) and  $\delta D = -726$  ‰ ( $\pm 20$  ‰). Our experimental values are in a very good agreement to the predicted value by Bottinga (1969) of  $\epsilon_{H_2-H_2O} = -722$  ‰, which gives a  $\delta D = -733$  ‰ (20 °C,  $\delta D$  of precipitation of  $-37.8$  ‰).

#### 4 Summary, conclusions and outlook

The isotopic composition of biologically produced H<sub>2</sub> was investigated systematically and our measurements confirm the massive deuterium depletion as predicted by Bottinga (1969). Using a Keeling plot analysis, we establish an overall source signature of  $\delta D = -712$  ‰ ( $\pm 13$  ‰) for biologically produced H<sub>2</sub>, with a fractionation constant of



**Fig. 2.** Dependency of H<sub>2</sub> mixing ratio (a) and ε<sub>H<sub>2</sub>-H<sub>2</sub>O</sub> (b) on incubation temperature for continuous incubations from the biogas plant under otherwise identical conditions (substrate 30 % grass, 30 % maize, 40 % cereals). Gray diamonds in figure (b) indicate the theoretically predicted value from Bottinga (1969). Results of the measured fit lines are: (a)  $y = 1990 \ln(x) - 7453$ ;  $R^2 = 0.96$ ; (b)  $y = 2.3x - 817.4$ ;  $R^2 = 0.98$ . Note: With respect to enzymatic catalysed production of hydrogen, a logarithmic fit is chosen for the relation between temperature and mixing ratio. Note: the error bars given are not representing the reproducibility of the duplicate measurements, but also take into account the general uncertainties by using mean relative errors of 2.9 % for δD and 5.4 % for the mixing ratio.

ε<sub>H<sub>2</sub>-H<sub>2</sub>O</sub> = 689 ‰ ± 20 ‰ between the H<sub>2</sub> and the source water at 38 °C and a δD<sub>H<sub>2</sub>O</sub> of -73.4 ‰. The temperature dependence of ε<sub>H<sub>2</sub>-H<sub>2</sub>O</sub> has also been determined, and accounting for the temperature effect the fractionation constant is extrapolated to ε<sub>H<sub>2</sub>-H<sub>2</sub>O</sub> = -731 ‰ (± 20 ‰) at 20 °C. This gives an experimentally received source signature of approximately δD = -741 ‰ (± 20 ‰) for biologically produced H<sub>2</sub> at mean temperatures (20 °C) and mean δD of precip-

itation (-37.8 ‰). Thus, we suggest using these values in global models, rather than the value of -628 ‰ that has been assumed in recent global model studies (Price et al., 2007; Pieterse et al., 2011).

As expected for a thermodynamic equilibrium, the isotopic fractionation is independent of used substrates in the samples from the biogas plant. Samples from individual microorganism cultures confirm the depletion in general, but show even slightly lower δD values; whereas H<sub>2</sub> produced from a nitrogen fixing species had slightly higher δD values. These differences could be caused by extremely high mixing ratios and dilution effects, but this needs further detailed investigation.

Due to its extreme deuterium depletion, biological H<sub>2</sub>, thus, has a high leverage in the global atmospheric H<sub>2</sub> isotope budget. Biological H<sub>2</sub> accounts for only ~ 10 % of the total H<sub>2</sub> source, but this fraction is depleted by ~ 772 ‰ relative to the ambient reservoir of ~ +130 ‰ (note that δ values do not add linearly), so including this source or not makes a huge difference of > 70 ‰ in the atmospheric isotope budget.

The new results imply that the δD values of biological H<sub>2</sub> are distinctly lower than what was included in the two recent global model studies of δD<sub>H<sub>2</sub></sub> (Price et al., 2007; Pieterse et al., 2011). First studies by Pieterse et al. (2011) included a sensitivity test for a change in the isotope source signature from -628 ‰ to -700 ‰ and found that this would change the atmospheric isotope budget by -4 ‰. The demand and production of biologically produced H<sub>2</sub> is expected to increase in the future, and a small increase in the production and release to the atmosphere of e.g., 1 Tg yr<sup>-1</sup> would lead to an observable decrease in δD in atmospheric H<sub>2</sub> and can influence the global isotope budgeting.

Despite the large advance in H<sub>2</sub> measurement techniques, the isotopic signature of this gas is still challenging to measure. Intercomparison experiments within the European “EUROHYDROS” project reveal mean deviations of < 1 % in H<sub>2</sub> and provide confidence in the reproducibility of the mixing ratios (Yver et al., 2011), but a lack of international isotope standards could still cause systematic errors in the isotopic scale. This should be taken into account when interpreting isotopic data.

**Acknowledgements.** We thank Susanne Voerkelius of Hydroisotop GmbH for the isotope analysis of the water samples in this project. Peter Haus, Tobias Mühlethaler, and Carolin Verbree have conducted the *Chlamydomonas reinhardtii* cultivation experiment as part of the middle school thesis project. Also thanks to Anton Hartmann, Research Center Munich, Germany for preparation of the bacillus *Azospirillum brasiliensis*.

This project was supported by the NWO (Netherlands Organisation for Scientific Research), NWO project number 816.01.001.

Edited by: U. Seibt

## References

- Batenburg, A. M., Walter, S., Pieterse, G., Levin, I., Schmidt, M., Jordan, A., Hammer, S., Yver, C., and Röckmann, T.: Temporal and spatial variability of the stable isotopic composition of atmospheric molecular hydrogen: observations at six EUROHYDROS stations, *Atmos. Chem. Phys.*, 11, 6985–6999, doi:10.5194/acp-11-6985-2011, 2011.
- Bottinga, Y.: Calculated fractionation factors for carbon and hydrogen isotope exchange in the system calcite-carbon dioxide-graphite-methane-hydrogen-water vapour, *Geochim. Cosmochim. Ac.*, 33, 49–64, 1969.
- Bowen, G. J. and J. Revenaugh: Interpolating the isotopic composition of modern meteoric precipitation, *Water Resour. Res.*, 39, 1299, doi:10.1029/2003WR002086, 2003.
- Ehhalt, D. H. and Rohrer, F.: The tropospheric cycle of H<sub>2</sub>: a critical review, *Tellus B*, 61, 500–535, 2009.
- Feck, T., Groß, J. U., and Riese, M.: Sensitivity of Arctic ozone loss to stratospheric H<sub>2</sub>O, *Geophys. Res. Lett.*, 35, L01803, doi:10.1029/2007GL031334, 2008.
- Feilberg, K. L., Johnson, M. S., Bacak, A., Röckmann, T., and Nielsen, C. J.: Relative tropospheric photolysis rates of HCHO and HCDO measured at the European photoreactor facility, *J. Phys. Chem. A*, 111, 9034–9046, 2007.
- Gerst, S. and Quay, P.: Deuterium component of the global molecular hydrogen cycle, *J. Geophys. Res.*, 106, 5021–5031, 2001.
- Hauglustaine, D. A. and Ehhalt, D. H.: A three-dimensional model of molecular hydrogen in the troposphere, *J. Geophys. Res.*, 107, 4330–4346, doi:10.1029/2001JD001156, 2002.
- Haus, P., Mühlethaler, T., and Verbree, C.: Wasserstoffproduktion mit Grünalgen, Maturarbeit, Kantonsschule Aarau, Switzerland, 2009.
- Hoffmann, G., Werner, M., and Heimann, M.: The water isotope module of the ECHAM atmospheric general circulation model – A study on time scales from days to several years, *J. Geophys. Res.*, 103, 16871–16896, 1998.
- Jacobson, M. Z.: Effects of wind-powered hydrogen fuel cell vehicles on stratospheric ozone and global climate, *Geophys. Res. Lett.*, 35, L19803, doi:10.1029/2008GL035102, 2008.
- Jacobson, M. Z., Colella, W. G., and Golden, D. M.: Cleaning the air and improving health with hydrogen fuel-cell vehicles, *Science*, 308, 1901–1905, 2005.
- Kaiser, J., Engel, A., Borchers, R., and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N<sub>2</sub>O isotope distribution, *Atmos. Chem. Phys.*, 6, 3535–3556, doi:10.5194/acp-6-3535-2006, 2006.
- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Worton, D. R., Sturges, W. T., Grunow, K., and Schmidt, U.: Contribution of very short-lived organic substances to stratospheric chlorine and bromine in the tropics – a case study, *Atmos. Chem. Phys.*, 8, 7325–7334, doi:10.5194/acp-8-7325-2008, 2008.
- Laube, J. C., Martinerie, P., Witrant, E., Blunier, T., Schwander, J., Brenninkmeijer, C. A. M., Schuck, T. J., Bolder, M., Röckmann, T., van der Veen, C., Bönisch, H., Engel, A., Mills, G. P., Newland, M. J., Oram, D. E., Reeves, C. E., and Sturges, W. T.: Accelerating growth of HFC-227ea (1,1,1,2,3,3,3-heptafluoropropane) in the atmosphere, *Atmos. Chem. Phys.*, 10, 5903–5910, doi:10.5194/acp-10-5903-2010, 2010.
- Nilsson, E. J. K., Johnson, M. S., Taketani, F., Matsumi, Y., Hurley, M. D., and Wallington, T. J.: Atmospheric deuterium fractionation: HCHO and HCDO yields in the CH<sub>2</sub>DO + O<sub>2</sub> reaction, *Atmos. Chem. Phys.*, 7, 5873–5881, doi:10.5194/acp-7-5873-2007, 2007.
- Nilsson, E. J. K., Andersen, V. F., Skov, H., and Johnson, M. S.: Pressure dependence of the deuterium isotope effect in the photolysis of formaldehyde by ultraviolet light, *Atmos. Chem. Phys.*, 10, 3455–3462, doi:10.5194/acp-10-3455-2010, 2010.
- Nimcevic, D., Schuster, M., and Gapes, J. R.: Solvent production by *Clostridium beijerinckii* NRRL B592 growing on different potato media, *Appl. Microbiol. Biot.*, 50, 426–428, 1998.
- Novelli, P. C., Lang, P. M., Masarie, K. A., Hurst, D. F., Myers, R., and Elkins, J. W.: Molecular hydrogen in the troposphere: Global distribution and budget, *J. Geophys. Res.*, 104, 30427–30444, 1999.
- Pieterse, G., Krol, M. C., and Röckmann, T.: A consistent molecular hydrogen isotope chemistry scheme based on an independent bond approximation, *Atmos. Chem. Phys.*, 9, 8503–8529, doi:10.5194/acp-9-8503-2009, 2009.
- Pieterse, G., Krol, M. C., Batenburg, A. M., Steele, L. P., Krummel, P. B., Langenfelds, R. L., and Röckmann, T.: Global modelling of H<sub>2</sub> mixing ratios and isotopic compositions with the TM5 model, *Atmos. Chem. Phys.*, 11, 7001–7026, doi:10.5194/acp-11-7001-2011, 2011.
- Prather, M. J.: An environmental experiment with H<sub>2</sub>?, *Science*, 302, 581–582, 2003.
- Price, H., Jaegle, L., Rice, A., Quay, P., Novelli, P. C., and Gammon, R.: Global budget of molecular hydrogen and its deuterium content: Constraints from ground station, cruise, and aircraft observations, *J. Geophys. Res.*, 112, D22108, doi:10.1029/2006JD008152, 2007.
- Rahn, T., Kitchen, N., and Eiler, J. M.: D/H ratios of atmospheric H<sub>2</sub> in urban air: Results using new methods for analysis of nanomolar H<sub>2</sub> samples, *Geochim. Cosmochim. Acta*, 66, 2475–2481, 2002.
- Rahn, T., Eiler, J. M., Boering, K. A., Wennberg, P. O., McCarthy, M. C., Tyler, S., Schauffler, S., Donnelly, S., and Atlas, E.: Extreme deuterium enrichment in stratospheric hydrogen and the global atmospheric budget of H<sub>2</sub>, *Nature*, 424, 918–921, 2003.
- Rhee, T. S., Mak, J., Röckmann, T., and Brenninkmeijer, C. A. M.: Continuous-flow isotope analysis of the deuterium/hydrogen ratio in atmospheric hydrogen, *Rapid Commun. Mass Sp.*, 18, 299–306, 2004.
- Rhee, T. S., Brenninkmeijer, C. A. M., and Röckmann, T.: The overwhelming role of soils in the global atmospheric hydrogen cycle, *Atmos. Chem. Phys.*, 6, 1611–1625, doi:10.5194/acp-6-1611-2006, 2006.
- Rice, A., Quay, P., Stutsman, J., Gammon, R., Price, H., and Jaegle, L.: Meridional distribution of molecular hydrogen and its deuterium content in the atmosphere, *J. Geophys. Res.*, 115, D12306, doi:10.1029/2009JD012529, 2010.
- Röckmann, T., Rhee, T. S., and Engel, A.: Heavy hydrogen in the stratosphere, *Atmos. Chem. Phys.*, 3, 2015–2023, doi:10.5194/acp-3-2015-2003, 2003.
- Röckmann, T., Gómez Álvarez, C. X., Walter, S., Veen, C. V., Wollny, A. G., Gunthe, S. S., Helas, G., Pöschl, U., Keppler, F., Greule, M., and Brand, W. A.: The isotopic composition of H<sub>2</sub> from wood burning - dependency on combustion efficiency,

- moisture content and  $\delta D$  of local precipitation, *J. Geophys. Res.*, 115, D17308, doi:10.1029/2009JD013188, 2010a.
- Röckmann, T., Walter, S., Bohn, B., Wegener, R., Spahn, H., Brauers, T., Tillmann, R., Schlosser, E., Koppmann, R., and Rohrer, F.: Isotope effect in the formation of H<sub>2</sub> from H<sub>2</sub>CO studied at the atmospheric simulation chamber SAPHIR, *Atmos. Chem. Phys.*, 10, 5343–5357, doi:10.5194/acp-10-5343-2010, 2010b.
- Schultz, M. G., Diehl, T., Brasseur, G. P., and Zittel, W.: Air pollution and climate-forcing impacts of a global hydrogen economy, *Science*, 302, 624–627, 2003.
- Sams, A. J. M., van Dijk, J. B., Dijkema, C., and Plugge, C. M.: Growth of syntrophic propionate-oxidizing bacteria with fumarate in the absence of methanogenic bacteria, *Appl. Environ. Microbiol.*, 59, 1114–1119, 1993.
- Sams, A. J. M. and Plugge, C. M.: Electron transfer in syntrophic communities of anaerobic bacteria and archaea, *Nat. Rev. Microbiol.*, 7, 568–577, doi:10.1038/nrmicro2166, 2009.
- Tromp, T. K., Shia, R.-L., Allen, M., Eiler, J. M., and Yung, Y. L.: Potential environmental impact of a hydrogen economy on the stratosphere, *Science*, 300, 1740–1742, 2003.
- Vollmer, M. K., Walter, S., Bond, S. W., Soltic, P., and Röckmann, T.: Molecular hydrogen (H<sub>2</sub>) emissions and their isotopic signatures (H/D) from a motor vehicle: implications on atmospheric H<sub>2</sub>, *Atmos. Chem. Phys.*, 10, 5707–5718, doi:10.5194/acp-10-5707-2010, 2010.
- Warwick, N. J., Bekki, S., Nisbet, E. G., and Pyle, J. A.: Impact of a hydrogen economy on the stratosphere and troposphere studied in a 2-D model, *Geophys. Res. Lett.*, 31, L05107, doi:10.1029/2003GL019224, 2004.
- Yver, C. E., Pison, I. C., Fortems-Cheiney, A., Schmidt, M., Chevalier, F., Ramonet, M., Jordan, A., Søvde, O. A., Engel, A., Fisher, R. E., Lowry, D., Nisbet, E. G., Levin, I., Hammer, S., Necki, J., Bartyzel, J., Reimann, S., Vollmer, M. K., Steinbacher, M., Aalto, T., Maione, M., Arduini, J., O'Doherty, S., Grant, A., Sturges, W. T., Forster, G. L., Lunder, C. R., Privalov, V., Paramonova, N., Werner, A., and Bousquet, P.: A new estimation of the recent tropospheric molecular hydrogen budget using atmospheric observations and variational inversion, *Atmos. Chem. Phys.*, 11, 3375–3392, doi:10.5194/acp-11-3375-2011, 2011.