



Responses of carbon dioxide flux and plant biomass to water table drawdown in a treed peatland in northern Alberta: a climate change perspective

T. M. Munir¹, B. Xu^{1,*}, M. Perkins¹, and M. Strack¹

¹Department of Geography, University of Calgary, Calgary, AB, Canada

* now at: NAIT Boreal Research Institute, Peace River, AB, Canada

Correspondence to: T. M. Munir (tmmunir@ucalgary.ca)

Received: 13 August 2013 – Published in Biogeosciences Discuss.: 13 September 2013

Revised: 10 December 2013 – Accepted: 3 January 2014 – Published: 11 February 2014

Abstract. Northern peatland ecosystems represent large carbon (C) stocks that are susceptible to changes such as accelerated mineralization due to water table lowering expected under a climate change scenario. During the growing seasons (1 May to 31 October) of 2011 and 2012 we monitored CO₂ fluxes and plant biomass along a microtopographic gradient (hummocks-hollows) in an undisturbed dry continental boreal treed bog (control) and a nearby site that was drained (drained) in 2001. Ten years of drainage in the bog significantly increased coverage of shrubs at hummocks and lichens at hollows. Considering measured hummock coverage and including tree incremental growth, we estimate that the control site was a sink of -92 in 2011 and -70 g C m⁻² in 2012, while the drained site was a source of 27 and 23 g C m⁻² over the same years. We infer that, drainage-induced changes in vegetation growth led to increased biomass to counteract a portion of soil carbon losses. These results suggest that spatial variability (microtopography) and changes in vegetation community in boreal peatlands will affect how these ecosystems respond to lowered water table potentially induced by climate change.

1 Introduction

Northern peatlands, functioning as carbon (C) sink ecosystems of the boreal forest over millennia, have stored approximately one third of global soil carbon (Tarnocai, 2006; Tarnocai et al., 2009; Turunen et al., 2002). These peatlands dominate the Canadian and Albertan landscape with cover-

age of 12 and 16 %, respectively and contain almost twice as much C per unit area (115 kg m⁻²) as tropical forests (Carlson et al., 2010; Vitt et al., 2009). Bogs in Western Canada (e.g. Alberta) are often covered by trees in contrast to open bogs in Eastern Canada (Turetsky et al., 2002). In Canada, the large peatland coverage (1.136 million km²) combined with high carbon density results in a store of approximately 147 Gt of soil organic C (Tarnocai, 2006). The large C pools formed as a result of net uptake of carbon dioxide (CO₂) from the atmosphere over millennia, if destabilized through a change in climate (e.g. atmospheric warming and subsequent drought), would lead to accelerated emission of greenhouse gases (GHGs) to the atmosphere (Gruber et al., 2004; IPCC, 2007; Limpens et al., 2008).

The formation and stability of peatland C stock is sensitive to changes in climatic conditions (e.g. atmospheric temperature and precipitation) (Vitt et al., 2009). Ongoing climate change is predicted to be most severe at northern latitudes where most of the peatlands are situated (Tarnocai, 2006; IPCC, 2007). The Canadian Global Climate Model (CGCM1, 2000) predicts a 3–4 °C increase in mean annual air temperature by 2020, with the greatest potential temperature increase (> 20 °C) occurring in winter months under extreme climate warming scenarios (Hengeveld, 2000). The increase of air temperature, combined with altered precipitation patterns, could lead to overall decrease in soil moisture across the high latitude region (IPCC, 2007). Drought/warming-induced water table drawdown could have a significant impact on the sustainability and ecosystem functions of boreal peatlands (Tarnocai, 2006; Adkinson et al.,

2011; Blodau and Siems, 2012; Chivers et al., 2009; Ise et al., 2008; Riutta, 2008, 2009; Roulet, 2000; Waddington and Price, 2000; Limpens et al., 2008). The increased atmospheric temperature and lowered water table can cause enhanced rates of organic matter decomposition and CO₂ emission, consequently resulting in the gradual depletion of peat C pool (Turetsky and Louis, 2006).

Carbon fluxes in peatlands occur in the forms of the uptake of C from the atmosphere via gross primary photosynthesis (GPP) and the release of C to the atmosphere by respiration (R) of plants (autotrophic) and microorganisms (heterotrophic). The sum of GPP and R is defined as the net ecosystem exchange (NEE) of CO₂. Net uptake of CO₂ causes the accumulation of carbon in the form of plant biomass and soil organic matter. The GPP, R and NEE of the forest floor are denoted by GPP_{ff}, R_{ff} and NE_{ff}, respectively.

Photosynthesis and autotrophic respiration may vary independently with changing temperature (Ryan, 1995; Ow et al., 2008). Warm and dry conditions in peatlands can either stimulate CO₂ uptake by enhanced GPP (e.g. Moore and Dalva, 1993; Updegraff et al., 2001; Syed et al., 2006; Ise et al., 2008; Cai et al., 2010) or reduce C uptake by limiting moisture (Alm et al., 1999; Roulet et al., 2007; Ise et al., 2008). Lowered water table in a treed bog increased tree productivity and fine root biomass significantly in a Canadian (Lieffers and Rothwell, 1987) and a Finnish peatland (Heikurainen and Pakarinen, 1982). Cool temperatures and near-surface water table conditions which typically occur in northern peatland ecosystems suppress respiration (Gorham, 1991; Hanson et al., 2000; Davidson and Janssens, 2006; Chapman and Thurlow, 1998). Predicted warming and subsequent lowered water table will result in enhanced CO₂ emissions from northern peatland soils (Moore, 2002; Roulet et al., 1992) where fine tree root biomass may contribute to soil total respiration (Lieffers and Rothwell, 1987). However, while the variation in respiration may not always be linked to fluctuation in water table, it is related to changes in moisture. Thus water table is an important control on respiration in peatlands in which soil peat moisture is sensitive to lowering of water table (Parmentier et al., 2009). Therefore, depending on the balance of GPP and R changes as a consequence of warming and/or drought, there may be a net increase or decrease in thickness of peat (Moore et al., 2006).

Autotrophic respiration by tree/shrub roots may contribute a significant amount to forest floor respiration (R_{ff}) when lowered water table stimulates root growth and promotes overall shrub/tree growth in dried peatlands (Lohila et al., 2011). Separating tree root respiration (R_r) from R_{ff} is critical in order to attribute the C losses to various sources of soil respiration and to better understand C source/sink dynamics (Hanson et al., 2000; Valentini et al., 2000; Janssens et al., 2001) of boreal treed peatlands in the face of global climate change. Isolating R_r from R_{ff} can make possible the comparison of CO₂ fluxes and plant biomass of a treed bog with those of an open bog, provided all controlling variables are

similar. The contribution of R_r to R_{ff} has been quantified using closed chamber technique in various forest ecosystems. Hermle et al. (2010) separated black spruce root respiration from soil total respiration by measuring the difference between control and trenched plots. They found that the R_r was 24 % of the soil total respiration. The contribution of R_r to soil total respiration was higher (37 %) in a subtropical forest (Wang et al., 2008) in a similar trenching experiment. An even higher contribution of rhizomicrobial respiration was quantified by Hanson et al. (2000) for forest vegetation in Florida.

As peatlands become drier under warming climate, it has been suggested that vegetation communities could shift towards a shrub/tree dominated system (Weltzin et al., 2000; Camill, 1999; Lohila et al., 2011), which in turn could alter the above (Lohila et al., 2011) and belowground C dynamics (Blodau and Siems, 2012). Swedish peatlands drained for forestry have been reported to respire at very high rates of 513–6516 g CO₂ m⁻² yr⁻¹ (von Arnold et al., 2005). A net loss of soil carbon in Finnish peatlands drained for forestry has also been reported with mean value of 150 g C m⁻² yr⁻¹ (Simola et al., 2012).

The shift in vegetation coverage and C dynamics vary with microtopographic features (e.g. hummocks (H) and hollows (W)) in peatlands (e.g. Strack et al., 2006, Waddington and Roulet, 2000). Also, the relationships between peatland CO₂ fluxes and water table may vary spatially between different microtopographic features in peatlands (Charman and Chichester, 2002; Joosten and Clarke, 2002). For example Strack et al. (2006) reported reduced GPP at hummocks and enhanced GPP at hollows and lawns in a water table draw-down experiment in an open, poor fen peatland. Bubier et al. (2003) reported a significant increase in total respiration at bog hollows during a dry summer and no change at hummocks.

Drought response experiments have been conducted in Eastern Canada (for example Waddington and Price, 2000; Strack et al., 2006) where generally most of the peatlands are characterized by their open nature (minimal tree cover) and receive high precipitation that leads to high surface humidity. Climatic and environmental (temperature, precipitation and water table position) response experiments have been conducted in Western Canada (Adkinson et al., 2011; Syed et al., 2006) where in contrast to Eastern Canada, most of the peatlands are generally drier and warmer and are characterized by their tree cover (Vitt et al., 1998; Price, 2010). However, these were short-term responses to drought studies and differences in microtopographic response were not considered. Western Canadian continental treed bogs are expected to respond to predicted climate change differently for CO₂ fluxes and plant biomass than those of Eastern Canadian open peatlands, with the potential for vegetation succession when water tables are persistently lowered. Moreover, we are unaware of any drought response CO₂ flux and biomass

change experiments in treed peatland that have measured contribution of R_r to R_{ff} using the closed chamber technique.

Therefore, our research aimed: (1) to compare CO_2 fluxes along a microtopographic gradient (hummock vs. hollow) between natural (control) and drained sites at a continental ombrotrophic bog, (2) to quantify changes in tree biomass and ground-layer biomass along microtopographic gradient in response to drainage, and (3) to determine the contribution of tree root respiration to forest floor respiration as affected by drainage.

2 Methods

2.1 Study sites

Two sections of a dry continental ombrotrophic bog were selected: one undisturbed section (CONTROL) (+55°21'14.19" N; -112°31'3.69" W) and one section that was drained in 2001 (DRAINED) (+55°16'44.28" N, -112°28'8.22" W). The drained site was not specifically drained for forestry but inadvertently drained during horticultural peat extraction operations on nearby sites. The drained site is located near the corner of two main ditches that have effectively drained a large quadrant of the peatland. All plots were within 50 m of the ditches. The two sections were 9 km apart and located in north-central Alberta, approximately 85 km northeast of Athabasca, Alberta, Canada. Both sites are underlain by sandy clay substrate and have peat depth exceeding 4 m. Climate in this region is sub-humid continental with mean annual and growing season (May–October) temperatures at 2.1 °C and 11.7 °C (Environment Canada, 2013). Mean annual precipitation at Athabasca is 504 mm, with 382 mm falling as rain. The research was conducted over two growing seasons (2011–2012). Mean growing season rainfall and air temperatures measured on site were 402.7 and 281.6 mm and 13.06 °C and 13.08 °C for 2011 and 2012, respectively.

The Wandering River, Alberta bog comes under the class of treed low shrub bogs, with typical mosaic of hummock (H) and hollow (W) microforms (Riley, 2003). The hummocks and hollows at the control site were dominated by *Sphagnum* mosses with sparse shrubs. The drained site had higher coverage of shrubs on the hummocks and higher lichen coverage in the hollows. Common mosses included *Sphagnum fuscum*, *Sphagnum magellanicum*, *Sphagnum capillifolium* and *Pleurozium schreberi* while common shrubs included Labrador Tea (*Rhododendron groenlandicum*), Lingonberry (*Vaccinium vitis-idaea*), small bog cranberry (*Oxycoccus microcarpus*) and cloudberry (*Rubus chamaemorus*). The most common tree in the bog was black spruce (*Picea mariana*) that constituted > 99 % of the tree stand with 25 766 stems ha^{-1} consisted of 37 % taller trees (> 137 cm height) up to 769 cm. The black spruce stand had an average canopy height of 168 cm, projection coverage of

42 % and basal area of 73.5 $\text{m}^2 \text{ha}^{-1}$. This description applies to the whole bog having control and drained sites.

2.2 CO_2 exchange

At each site, three hummocks and three hollows were chosen as the study plots before the growing season (May–October) of 2011. A 60 cm × 60 cm steel collar having grooves at the top was inserted about 6 cm into the peat at each plot to keep disturbance to roots minimal. The CO_2 fluxes were measured weekly during the growing seasons using a closed chamber having dimensions of 60 cm × 60 cm × 30 cm ($L \times D \times H$), made of clear acrylic and corrected for transmittance (88 %). Two small battery-operated fans were installed inside the chamber to circulate the air and achieve equilibrium CO_2 concentration between measurements. The instantaneous CO_2 concentration in the chamber was monitored with a portable infrared gas analyser (PP systems, USA, EGM-4). Photosynthetically active radiation (PAR) was measured with a quantum sensor (PP systems, USA) placed at the top of the chamber. The temperature inside the chamber was measured with a thermocouple thermometer (VWR Int., USA). All measurements were made at 15 s intervals for up to 1.75 min. At the time of flux measurements, soil temperatures at the depths of 2, 5, 10, 15, and 20 cm were measured with a thermocouple thermometer at all plots. Water table relative to moss surface was measured at each CO_2 flux measurement occasion from a permanently installed well associated with each plot. We used negative values for indicating belowground water table.

At each plot, a total of 184 CO_2 flux measurements were made during the daytime of growing seasons (May–October) of 2011 and 2012. During the two growing seasons we organized 20 flux measurement campaigns. Each campaign lasted for about eight days during which fluxes were measured on two to three occasions at each plot. At each flux measurement occasion we took a total of five measurements (four measurements for NE_{ff} (net exchange of CO_2 at the forest floor) under a range of PAR levels created using shades, and a last run for R_{ff}) each of 1.75 min: 2–3 full sun, single shade, double shade and finally opaque tarp (for R_{ff}). The chamber was flushed (vented) for enough time between the measurements to bring the headspace concentration in equilibrium with ambient air concentration. Therefore we measured respiration as the final measurement (after about 18 min) at each plot by using the clear chamber covered with an opaque shroud. In this way any buildup of CO_2 in the soil would have already been flushed. Thus problems in determining respiration rates caused by flushing CO_2 built up in the soil during night time chamber measurements (Lai et al., 2012; Koskinen et al., 2013) were not an issue for our measurements.

As noted above, CO_2 flux measurements in the dark (when the chamber was covered with an opaque shroud) represented R_{ff} . We recognize that this R_{ff} represents only forest floor respiration including understory aboveground biomass

Table 1. Estimated parameter values (\pm SE), standard error of the estimate (SEE) and goodness of fit (r^2) for the gross primary production of forest floor (GPP_{ff}) Model (Eq. 1)*.

Year	Site/Microform	GPP _{ff} vs. PAR		GPP _{ff} vs. WT		GPP _{ff} vs. <i>T</i>		SEE (g CO ₂ m ⁻² d ⁻¹)	<i>r</i> ²
		<i>P</i> _{max} (g CO ₂ m ⁻² d ⁻¹)	<i>k</i> (μmol m ⁻² s ⁻¹)	WT _{opt} (cm)	WT _{tol} (cm)	<i>T</i> _{opt} (°C)	<i>T</i> _{tol} (°C)		
2011	Control Hummocks	-22.2 ± 4.6	900 ± 422	-56.0 ± 13.0	30.0 ± 15.8	16.1 ± 3.7	10.0 ± 5.4	2.30	0.83
	Control Hollows	-19.4 ± 7.1	950 ± 601	-28.3 ± 9.7	22.9 ± 21.3	14.1 ± 7.5	10.0 ± 12.5	2.05	0.63
	Drained Hummocks	-35.9 ± 8.8	950 ± 440	-118.1 ± 8.3	30.0 ± 11.1	12.3 ± 1.5	10.0 ± 3.1	1.14	0.82
	Drained Hollows	-29.2 ± 16.8	850 ± 639	-70.5 ± 31.6	30.0 ± 17.3	10.6 ± 2.7	10.0 ± 10.2	0.57	0.68
2012	Control Hummocks	-24.8 ± 4.7	900 ± 286	-34.9 ± 11.3	30.0 ± 14.0	15.2 ± 2.0	10.0 ± 3.8	1.84	0.86
	Control Hollows	-21.8 ± 3.9	950 ± 299	-42.3 ± 13.8	30.0 ± 20.1	14.6 ± 1.3	9.9 ± 2.1	1.09	0.85
	Drained Hummocks	-50.0 ± 14.3	950 ± 416	-104.8 ± 2.8	30.0 ± 2.9	10.1 ± 2.0	10.0 ± 2.0	0.96	0.82
	Drained Hollows	-31.3 ± 10.1	850 ± 246	-99.5 ± 11.1	23.2 ± 6.7	10.0 ± 5.5	9.9 ± 4.6	1.55	0.83

* The models were developed for each microform type ($n = 3$) at the control and drained sites separately for growing seasons of 2011 and 2012. PAR, WT and *T* represent photosynthetically active radiation, water table (negative values show belowground water level) and 5 cm soil temperature, respectively. *P*_{max} denotes potential maximum rate of GPP_{ff} and *k* is level of PAR at which half of GPP_{ff} occurs. WT_{opt} and *T*_{opt} are Gaussian response parameters for optimum GPP_{ff} while WT_{tol} and *T*_{tol} are Gaussian response parameters describing the width of the curve. All modeled parameters are significant at $p = 0.05$ level.

respiration, heterotrophic soil respiration and tree root respiration, and ignores respiration of the overstory aboveground tree biomass. GPP_{ff} was determined as the difference between NE_{ff} and *R*_{ff}. We used the convention that negative values indicate an uptake of CO₂ by the ecosystem.

Net exchange of CO₂ at the forest floor (NE_{ff}) was calculated using exponential change (Kutzbach et al., 2007) instead of linear change in CO₂ concentration in the chamber headspace with time, as a function of volume, air temperature and pressure inside the chamber, according to ideal gas law. The exponential regression was used because covering soil and/or vegetation essentially manipulates the spontaneous CO₂ fluxes by altering the concentration gradients between the soil, the vegetation and the air inside the chamber. Due to the constantly changing controls on CO₂ flux within the chamber, no linear decrease or increase of CO₂ concentration inside the chamber can be expected. Kutzbach et al. (2007) found that the linear CO₂ fluxes compared with exponential fluxes were up to 40 % lower, over CO₂ chamber closure time of only two minutes.

Maximum rates of GPP_{ff} (GPP_{max}) and NE_{ff} (NE_{max}) represent GPP_{ff} and NE_{ff} when the photon flux density of photosynthetically active radiation is greater than 1000 μmol m⁻² s⁻¹. As modeled values of the maximum rate of photosynthesis are likely never achieved in reality, these values represent a more realistic estimate of CO₂ exchange when light is not limiting as discussed in Bubier et al. (2003). We use these to statistically compare between plots to better understand processes (changes in plant cover, species type, water table, etc.) that affect CO₂ exchange. Data reported are averages for the study seasons (May–October) for all occasions when PAR > 1000 μmol m⁻² s⁻¹.

2.2.1 Seasonal CO₂ exchange model

The growing season (1 May to 31 October) GPP_{ff} was estimated using an empirical model following Riutta et al. (2007) parameterized separately for each microform type × water table treatment. The parametrization was done separately for each of the growing seasons of 2011 and 2012. The seasonal GPP_{ff} was estimated by

$$\text{GPP}_{\text{ff}} = \frac{\text{PAR} \times P_{\text{max}}}{\text{PAR} + k} \times e^{\left[-0.5 \times \left(\frac{\text{WT} - \text{WT}_{\text{opt}}}{\text{WT}_{\text{tol}}}\right)^2\right]} \times e^{\left[-0.5 \times \left(\frac{T - T_{\text{opt}}}{T_{\text{tol}}}\right)^2\right]}, \quad (1)$$

where *P*_{max} denotes the potential maximum rate of GPP_{ff} (g CO₂ m⁻² d⁻¹) if water table and temperature are not limiting and the parameter *k* denotes the level of PAR at which half of the *P*_{max} occurs. WT is the water table position (cm), WT_{opt} and WT_{tol} are parameters in a Gaussian response of GPP_{ff} to water table when GPP_{ff} is optimized and width of the curve respectively, *T* is the soil temperature (°C) at 5 cm below moss surface and *T*_{opt} and *T*_{tol} are parameters in a Gaussian response of GPP_{ff} to the soil temperature when GPP_{ff} is optimized and width of the curve.

Model parameters for seasonal GPP_{ff}, standard errors (\pm), r^2 values, and standard errors of the estimates at control and drained microforms are presented in Table 1. Two-thirds of the data were randomly selected and used for model construction, whereas one-third of the data were used for independent testing of the models following Tuittila et al. (2004).

After examining the data it appeared that the relationship of *R*_{ff} with soil temperature at 5 cm depth was not exponential. Therefore the growing season *R*_{ff} was estimated using multiple linear regressions with soil temperature at 5 cm

Table 2. Estimated regression coefficient values (\pm SE), standard error of the estimate (SEE) and goodness of fit (r^2) for the forest floor respiration (R_{ff}) model (Eq. 2)*.

Year	Site/Microform	a	b	c	SEE (g CO ₂ m ⁻² d ⁻¹)	r^2
2011	Control Hummocks	1.30 \pm 0.29	-0.21 \pm 0.12	-18.25 \pm 6.80	0.80	0.81
	Control Hollows	1.70 \pm 0.32	-0.61 \pm 0.20	-32.33 \pm 10.74	0.68	0.79
	Drained Hummocks	0.31 \pm 0.27	-0.02 \pm 0.03	2.62 \pm 7.26	0.38	0.49
	Drained Hollows	0.55 \pm 0.13	-0.02 \pm 0.03	2.26 \pm 3.63	0.19	0.85
2012	Control Hummocks	0.39 \pm 0.15	-0.25 \pm 0.09	-10.71 \pm 5.31	0.61	0.53
	Control Hollows	0.93 \pm 0.08	-0.25 \pm 0.05	-12.77 \pm 1.77	0.36	0.81
	Drained Hummocks	0.65 \pm 0.06	-0.03 \pm 0.05	-3.40 \pm 8.14	0.32	0.91
	Drained Hollows	1.07 \pm 0.09	-0.15 \pm 0.04	-17.31 \pm 5.55	0.86	0.80

* The models were developed for each microform type ($n = 3$) at the control and drained sites separately for growing seasons of 2011 and 2012; a , b and c are regression coefficients. Negative values of b represent greater respiration with deeper water table values (belowground WT having negative values). All modeled parameters are significant at $p = 0.05$ level.

depth and water table position by

$$R_{ff} = a \times T + b \times WT + c, \quad (2)$$

where a , b and c are regression coefficients (Table 2).

Seasonal GPP_{ff} and R_{ff} were estimated based on Eqs. (1) and (2) for each twenty minute period between 1 May and 31 October, averaged daily and summed for a growing season total based on measurements made on sites for PAR (LI-190, LI-COR, Nevada, USA), WT (Levellogger Junior, Solinst, USA) and temperature (Onset HOBOWare Pro, MA, USA). Growing season ground-layer NE_{ff} was determined by adding seasonal GPP_{ff} to seasonal R_{ff} estimates. The control site was instrumented with one additional soil temperature sensor (T 109, Campbell Scientific Inc., Utah, USA; depth = 5 cm) and one tipping bucket rain gauge (TE 525, Campbell Scientific Inc., Utah, USA; height = 150 cm) both wired to a data logger (CR 1000, Campbell Scientific Inc., Utah, USA) programmed to measure every minute and record the average at 20 min intervals.

2.2.2 Model validation

Of all the data measured in the field, we separated one third randomly and did not use it for model construction. The remaining two-thirds of the data were used for model construction. The unused data were later correlated with modeled data for validation of the GPP_{ff} and R_{ff} models. Validation of the model showed excellent agreement between predicted and measured values (Fig. 1a and b).

2.3 Tree root respiration

To exclude the contribution of R_r to R_{ff} , a trenching method (Wang et al., 2008) was used. In May 2012, a total of 32 plots (eight hummocks and eight hollows at each site, in addition to the already described CO₂ flux plots) were chosen randomly from the available microtopography. At each site four hummock and four hollow plots of area 60 cm \times 60 cm were

cut around up to approximately 30 cm depth in three intervals (0–10 cm, 10–20 cm, 20–30 cm). It was assumed that cutting the peat would cut down most of the live root ingrowth. To make the cut loose enough to insert polyethylene sheet, we had to use a saw several times at each of the three depth intervals. Then the polyethylene sheet was inserted deep to 30 cm to prevent root ingrowth and the cuts were infilled with soil in the reverse order of removal (i.e., first we filled back the soil from 20–30 cm depth followed by 10–20 cm and lastly 0–10 cm). Although this procedure did not ensure that the backfilled soil occupied its original place, our intention was to keep disturbance minimal. The remaining four hummocks and four hollows were left intact to quantify the difference in CO₂ emission between cut (having minimal tree roots) and intact (with all tree roots) plots. During July to September 2012, all plots were clipped every two weeks to ensure that soil surface was free of live mosses, shrubs and herbs following Hanson et al. (2000), Riutta et al. (2007) and Hermle et al. (2010). The trenched and intact plots were clipped so that we could isolate soil respiration (measured at trenched plots) from R_r + soil respiration (measured at intact plots). Had the plots not been clipped, we would have measured R_r + soil respiration + autotrophic respiration of surface vegetation at intact plots and soil respiration + autotrophic respiration of surface vegetation at trenched plots. This way we could not have R_r separated from soil respiration. The surface vegetation was clipped with Fiskars power lever shears (Model # 100017192) that clips horizontally to keep disturbance minimal.

The CO₂ emissions from all plots were measured using the same instruments and chamber used for the measurement of NE_{ff} and R_{ff} and hence GPP_{ff}. We had a methodological challenge that while the cutting separates R_r from R_{ff} , it also adds fresh litter to the soil that can add to the existing heterotrophic soil respiration.

Trenching or cutting experiments have been performed to separate root autotrophic respiration from R_{ff} (e.g. Hanson




Fig. 1. Goodness of fit (r^2) between (a) modeled and measured GPP_{ff} and (b) modeled and measured R_{ff} . The measured data used in these figures were chosen randomly and were not used for construction of models. Both 2011 and 2012 data from control and drained sites are presented. Lines represent the 1 : 1 fit.

et al., 2000; Hermle et al., 2010; Wang et al., 2008; Díaz Pinés et al., 2010; Kuzyakov, 2006; Brant et al., 2006). In all cases the assumption has been made that the trenched roots die off within a short time and that afterwards the measured R_{ff} can solely be attributed to heterotrophic soil respiration. Trenching immediately disrupts the supply of recent photosynthates to the roots and mycorrhiza. The mycorrhizal fungi and associated bacteria will suffer from the lack of labile C. Bowden et al. (1993), Boone et al. (1998) and Rey et al. (2002) in trenching or cutting experiments showed that C content of decomposing fine roots in trenched plots contribute little to R_r and become stable in the four months after trenching. Therefore, no correction for extra CO_2 from decaying fine roots is necessary. However, the root exclusion experiment may not be useful if extended through a complete annual cycle, as over such a long period there is the possibility of reinvasion of roots into the previously root-free trenched plot (Edwards and Norby, 1999). While it is clear that findings from such trenching measurements should be interpreted carefully, the primary focus of this paper is to quantify R_{ff} while investigating R_r to better understand and separate the contribution of various processes to shifts in R_{ff} following drainage.

2.4 Biomass and tree productivity

Forest floor biomass was measured by clipping 25 cm × 25 cm quadrats at nine hummocks and nine hollows, at each of control and drained sites, in mid-July 2011. The biomass was clipped at the base of the capitulum at 1.0 cm below moss surface following Clymo (1970) and Loisel et al. (2012). From triplicate of each microform at each site, soil cores of only 20 cm depth were collected

due to frozen peat beyond this depth at the time of sample collection. The soil cores were sectioned into two depths (0–10 and 10–20 cm) and roots were sorted into coarse (> 2 mm) and fine (< 2 mm) fractions. For tree biomass, we selected three 10 m × 10 m quadrats in areas directly surrounding the flux plots at each site. The total study areas were not large and these plots covered most of the trees in the study areas. Trees were divided into tall (> 137 cm height) and short (< 137 cm height) for biomass estimation. All trees were measured for their height, diameter at breast height (DBH, when tall enough) and basal diameter (DB). Tall tree biomass was calculated by using an allometric equation (dry biomass = 0.153(tree DBH)^{2.248}) from Grigal and Kernik (1984). Trees < 137 cm were not measured for DBH as their total height was below a standard DBH measurement height. A subsample of 20 smaller trees > 125 cm were harvested parallel to the forest floor and taken back to the lab and oven dried at 80°C for 48 hours. The height and dry biomass of each tree was measured and an exponential regression was performed. An allometric equation was generated by regressing height with oven-dried weight as dry biomass = 0.0085(tree height)^{2.2088} ($R^2 = 0.93$, $p < 0.001$).

For calculating NPP of the tree stand, we adapted methods of Szumigalski and Bayley (1996) and Thormann and Bayley (1997). In addition to fens, they also estimated NPP of an Alberta ombrotrophic treed bog of hummock-hollow microtopography by adding aboveground incremental biomass to stand litter production (17 % of incremental biomass $m^{-2} yr^{-1}$ for *Picea mariana*). We quantified the incremental biomass of tall trees for 2011 and 2012 based on tree ring widths using DendroScan (Varem-Sanders et al., 1996). The incremental biomass of short trees for 2011 and 2012 was calculated by regressing leader length with height following

Table 3. Ground-layer and tree biomass (g m^{-2}) at control and drained sites.

Site/ Microform	ABOVEGROUND				BELOWGROUND			TREES	TOTAL
	Moss	Vascular	Lichen	Total	Fine (< 2 mm)	Coarse (> 2 mm)	Total		
Control Hummocks	215 ± 68 ^A	218 ± 84 ^A	0 ± 0 ^A	433 ± 147 ^A	563 ± 103	138 ± 31	701 ± 106	2142 ± 376	3304 ± 793
Control Hollows	182 ± 93 ^A	135 ± 41 ^A	11 ± 32 ^A	328 ± 72 ^A	484 ± 195	386 ± 425	870 ± 553		
Drained Hummocks	16 ± 27 ^B	737 ± 60 ^B	13 ± 14 ^A	766 ± 323 ^B	470 ± 96	498 ± 262	967 ± 275	1964 ± 381	3676 ± 1029
Drained Hollows	80 ± 130 ^A	133 ± 44 ^A	358 ± 354 ^B	571 ± 280 ^{AB}	491 ± 60	626 ± 392	1118 ± 422		

* Values are means ± SE ($n = 9$ for aboveground ground layer; $n = 3$ for belowground; $n = 3$ for trees). Means sharing same letters do not differ significantly. Letters should be compared only within one column. Total biomass was determined by weighting ground layer by the proportion of hummocks and hollows at each site (control = 56 % hummocks; drained = 52 % hummocks).

Mullin et al. (1992) and Macdonald and Lieffers (1990). Summation of biomass increments of tall and short trees for a year represented incremental biomass of tree stand for aboveground parts of the trees ($\text{IC}_{\text{biom_ag}}$) for that year at either site. However, litterfall was not estimated. Therefore we predicted a value of stand litter production based on Szumigalski and Bayley (1996; 17 % of incremental biomass) for Alberta ombrotrophic bogs. We also did not measure incremental biomass of the belowground parts of the tree ($\text{IC}_{\text{biom_bg}}$) due to the difficulty in measuring this component without disturbing our study sites for future monitoring. Therefore, we use an allometric equation (tree root biomass = 0.222* tree aboveground biomass) from Li et al. (2003) for estimating incremental biomass of tree roots.

2.5 CO₂-C balance calculations

The CO₂-C balances of the treed control and drained sites (NEE) were calculated separately for the growing seasons (1 May to 31 October) of 2011 and 2012 as

$$\text{NEE} = \text{NE}_{\text{ff}} + \text{IC}_{\text{tree_ag}} + \text{IC}_{\text{tree_bg}} + L_{\text{tree}} - R_{\text{r}}, \quad (3)$$

where NEE denotes net ecosystem exchange, NE_{ff} represents net exchange of CO₂-C of the forest floor, $\text{IC}_{\text{tree_ag}}$ and $\text{IC}_{\text{tree_bg}}$ represent incremental biomass growth of the aboveground and belowground parts of the trees, respectively, L_{tree} is tree litter production and R_{r} is tree root respiration.

Seasonal CO₂ fluxes at hummocks and hollows were up-scaled by multiplying mean estimated growing season CO₂ exchange by their respective coverage of 56 and 44 % and 52 and 48 % at the control and drained sites, respectively (Table 4). The incremental growth of the trees and their litter production was added to the forest floor CO₂ exchange assuming that biomass had a carbon content of 50 %. R_{r} was excluded to avoid double counting while determining CO₂-C balance of sites as $\text{IC}_{\text{tree_bg}}$ already accounts for R_{r} . We estimated a seasonal value of R_{r} by determining it as a proportion of R_{ff} based on instantaneous measurements and then estimating it as this proportion of the modelled seasonal R_{ff} .

2.6 Statistical analysis

Differences in GPP_{max} , R_{ff} , NE_{max} , and aboveground biomass between sites and microforms were tested by two-way ANOVA, using Minitab 16.0 (Minitab Inc., PA, USA). Differences in R_{r} between sites, microforms and cut and intact plots were tested for significance employing a three-way ANOVA using SPSS 20.1. The nonlinear and linear regression models (Eqs. 1, 2 and 3) were used to construct GPP_{ff} and R_{ff} models (SPSS 20.1) and to estimate seasonal CO₂-C balance.

3 Results

3.1 Site conditions

Ten years after initial drainage, the water table at the drained site was as much as 80 cm lower than that at the control site (Fig. 2). The growing seasons of 2011 and 2012 were warmer by 1.36 and 1.38 °C, respectively, and wetter by 41.9 mm in 2011 and drier by 79.2 mm in 2012 than 30 yr average at Athabasca. In 2012, the reduction in rainfall by 121 mm led to a decrease in water table level at control and drained hollows by 4.5 and 4.3 cm and at control and drained hummocks by 8.0 and 7.2 cm, respectively (Fig. 2).

The drained site was ditched around in 2001 and the data on pre-drawdown hydrology and vegetation were not available. However, given that the control and drained sites were part of the same bog and had similar vegetation layers (canopy layer consisted of *Picea mariana* and ground layer consisted of similar shrubs and mosses), air temperature, peat depth and underlain substrate are assumed to be similar before start of this study in 2001. As a result of 10 yr drainage, *Sphagnum* coverage at the drained site was significantly reduced by 97 % ($F(3, 32) = 33.40$, $p < 0.001$) compared to the control site, but no significant difference in *Sphagnum* coverage was observed between microforms at either site (Fig. 3). *Sphagnum* at drained site was replaced by shrubs at hummocks and lichens at hollows (field observation, data not presented here). The significant reduction in coverage of *Sphagnum* at the drained site was due to the unfavourable condition of very low water potential in the surface soil.

Table 4. Growing season CO₂-C flux estimates (\pm SE, g C m⁻²)^a.

Year	Site	GPP _{ff}	R _{ff}	NE _{ff}	R _r	L _{tree}	IC _{tree_ag}	IC _{tree_bg}	NEE ^b
2011	CONTROL								
	Hummock	-190 ± 29	225 ± 18	35 ± 30	63 ± 5.1	-6 ± 1.4	-38 ± 7.9	-6 ± 1.4	-92 ± 11.7
	Hollow	-178 ± 29	130 ± 21	-48 ± 9	12 ± 0.3				
	DRAINED								
	Hummock	-280 ± 21	295 ± 10	15 ± 7	62 ± 4.9	-11 ± 1.4	-66 ± 9.1	-15 ± 1.7	27 ± 13.6
	Hollow	-116 ± 19	536 ± 9	420 ± 23	123 ± 0.6				
2012	CONTROL								
	Hummock	-228 ± 43	216 ± 33	-12 ± 11	60 ± 5.1	-6 ± 1.2	-33 ± 6.8	-7 ± 1.2	-70 ± 10.2
	Hollow	-181 ± 8	241 ± 45	60 ± 33	21 ± 0.3				
	DRAINED								
	Hummock	-333 ± 75	359 ± 18	26 ± 13	75 ± 4.9	-10 ± 1.8	-60 ± 10.3	-13 ± 1.9	23 ± 14.9
	Hollow	-118 ± 20	507 ± 35	389 ± 47	116 ± 0.6				

^a Negative values represent uptake of carbon by the peatland from the atmosphere. The forest floor respiration (R_{ff}) includes tree root respiration (R_r). ^b NEE is calculated using equation 3 ($NEE = NE_{ff} + IC_{tree_ag} + IC_{tree_bg} + L_{tree} - R_r$). Forest floor carbon exchange was determined by weighting NE_{ff} measured at each microform by the proportion of hummocks and hollows at each site (control = 56 % hummocks; drained = 52 % hummocks).

3.2 Biomass and incremental tree growth

Vascular plant biomass at the drained hummocks was significantly higher than that at the control hummocks ($F(1, 32) = 17.07$, $p < 0.001$) and there was a significant interaction between drainage and microform ($F(1, 32) = 35.74$, $p < 0.001$), while there was no difference between control and drained hollows (Table 3, Fig. 3). Conversely, moss biomass at drained hummocks was significantly lowest of all plots ($F(1, 32) = 26.28$, $p < 0.001$). In fact, moss biomass at the drained site was overall much lower than at the control site regardless of microform type, indicating a strong decline of moss cover with drainage. Lichen biomass on the other hand showed an increase following drainage, but it was the drained hollows that had the highest lichen biomass (over 30 times higher than control hollows; $F(1, 32) = 7.9$, $p = 0.008$) and the interaction between drainage and microform was statistically significant. As a whole, above-ground biomass was highest at drained hummocks ($F(1, 32) = 14.24$, $p = 0.003$) while lowest at control hollows. Neither total belowground root biomass nor tree biomass were significantly different between microforms and/or sites. However, total root biomass was higher in the drained site than that in the control (Table 3).

Although tree biomass was higher in the control site by 178 g m⁻², the annual aboveground tree increment during the study years (2011 and 2012) was significantly higher in the drained site (66 and 60 g C m⁻²) than the control (38 and 33 g C m⁻²) ($F(1, 3) = 3025$, $p = 0.012$). Using equations presented in Li et al. (2003) belowground tree increment was estimated as 6 and 7 g C m⁻² at the control site and 15 and 13 g C m⁻² at the drained site in 2011 and 2012, respectively (Table 4).

Spatial variability in tree stands is a generic characteristic of natural/peatland ecosystems and we did not have tree

stand data prior to the study period. Heterogeneity even between the three quadrats constructed at each site was large; however, the size of the study areas precluded our ability to include more replicates. Therefore, while we cannot be certain that the biomass was identical before the study, they were likely similar and we did see a clear change in tree growth (based on the tree rings) coinciding with the ditching 10 yr ago. Thus we are confident that the changes in incremental growth determined represent a clear response to the changing water table.

3.3 CO₂ fluxes

3.3.1 Measured CO₂ Fluxes

Drainage did not change GPP_{max} significantly in 2011 or 2012 (Fig. 4; two-way ANOVA, 2011: $F = 0.06$, $p = 0.813$, 2012: $F = 4.13$, $p = 0.08$). However, GPP_{max} was significantly higher at hummocks than hollows (2011: $F = 7.84$, $p = 0.027$, 2012: $F = 8.99$, $p = 0.017$). Drainage had a significant interaction with microtopography in 2012 leading to significantly different GPP_{max} at drained microforms. Drainage resulted in significantly higher R_{ff} (2011: $F = 6.85$, $p = 0.037$, 2012: $F = 8.52$, $p = 0.019$), but R_{ff} remained statistically similar between microforms at both sites in both years. The drained hollows were the largest sources of CO₂ emission largely due to the significantly higher contribution of R_r (5.03 g CO₂ m⁻² d⁻¹) to R_{ff} (18.02 g CO₂ m⁻² d⁻¹) than that of the R_r contribution of 1.51 g CO₂ m⁻² d⁻¹ to R_{ff} (11.84 g CO₂ m⁻² d⁻¹) at control hollows (see 2012 in Fig. 4). NE_{max} was positive in 2011 but became negative (net sink of CO₂) in 2012 at control microforms. Subtracting R_r from NE_{max} switched the drained hummocks to a moderate sink and the control microforms to




Fig. 2. Water table levels (lines) and daily precipitation (bars) during growing seasons of 2011 and 2012. The cumulative seasonal precipitation during 2012 was 30 % less than that of 2011.




Fig. 3. Relationship between drainage/microform scenario and ground-layer aboveground biomass by category as a percentage of total.

larger sinks of CO₂, while considerably reduced emissions at the drained hollows (Fig. 4).

3.3.2 Modeled CO₂ fluxes

Based on empirical models (Eqs. 1 and 2), the ground layer at the control site was a small growing season sink of CO₂ taking up an estimated 6.9 g CO₂ m⁻² largely due to the significantly higher GPP_{ff} at its hollows than that of its hummocks. In contrast, the ground layer at the drained site was a substantial source of CO₂, losing an estimated 770 g CO₂ m⁻² largely due to significantly higher R_{ff} at its hollows than that of its hummocks, during the 2011 growing season (Table 4). In 2012, a shift in the functions of hollows and hummocks

at the control site was noticed, where hummocks became a moderate sink of CO₂, and the hollows became a substantial source. However, the drained microforms and site remained consistently sources of CO₂ (Table 4).

3.4 CO₂-C balance

In the growing season of 2011, the forest floor (including R_f) of control and drained sites were a small sink (2 g CO₂-C m⁻²) and substantial source (210 g CO₂-C m⁻²), respectively. In the growing season of 2012, the control site became a moderate source (20 g CO₂-C m⁻²) while the drained site remained a substantial source (200 g CO₂-C m⁻²). To calculate the final CO₂-C balance, we added estimated NE_{ff} (-R_f)




Fig. 4. GPP_{max}, R_{ff} and NE_{max} at control and drained sites in 2011. The R_r measurements were added in 2012. NE_{max} (without above- and belowground parts of trees) was calculated by subtracting R_r from NE_{max} and represents net exchange of CO₂ of the ground-layer vegetation and peat. H and W represent hummocks and hollows, respectively. Error bars indicate \pm standard deviation. Results are from two-way (drainage and microform) ANOVA performed separately for growing season of each year. Differences were evaluated between study plots grouped according to microform and drainage and are indicated by letters at each bar. Sites are significantly different at $p < 0.05$ if they have no letters in common (letters should be compared only within one CO₂ component in one year).

to estimated tree incremental growth ($IC_{tree_ag} + IC_{tree_bg}$) and tree litter production (L_{tree}) during the study years, and estimated that the control site was a larger sink of -92 g C m^{-2} in 2011 than that of -70 g C m^{-2} in 2012. However, the drained site remained a source through both study seasons, losing 27 g C m^{-2} in 2011 and 23 g C m^{-2} in 2012 growing season. Importantly, majority of the discussion in this study focuses on the differences in the measured components as opposed to the CO₂-C balance itself.

4 Discussion

The control site of this sub-humid, continental treed bog was a growing season sink of CO₂-C of 92 and 70 g C m^{-2} in years slightly wetter and drier than average, respectively. Depending on time since fire, Wieder et al. (2009) report that treed bogs in the same region represent an annual CO₂ sink of 120 to 220 g C m^{-2} and thus our value is slightly below this range. Within the same region of northern Alberta as the present study, Adkinson et al. (2011) report net growing season CO₂ exchange across three study years of -110.1 and -153.5 to -34.5 g C m^{-2} at poor fen and rich fen sites, respectively.

Previous research has shown that warm and dry summer conditions can reduce net CO₂ uptake in peatlands by enhancing respiration greater than productivity (Alm et al., 1999; Arneth et al., 2002; Bubier et al., 2003; Aurela et al., 2007). Similarly, in our experiment drier weather in 2012 reduced net uptake of CO₂ and reduced the growing season C sink at the control site. The shift was due to the substantially increased R_{ff} at the hollows greater than that of combined

increase in GPP_{ff} at the microforms (Table 4). The enhanced R_{ff} at hollows might be due to stressed vegetation growth observed at the drier hollows (Fig. 3). In contrast there was little change in GPP_{ff} or R_{ff} at the drained site in 2012 and thus no real change in net CO₂ emission.

Ten years of water table drawdown, converted our bog site forest floor from a small sink of 2 g m^{-2} in 2011 or a smaller source of 20 g m^{-2} in 2012 to a large source of CO₂ of $\sim 200 \text{ g m}^{-2}$ in both years. This value of net source of CO₂ compares well with those of other drained peatlands as reported by Waddington et al. (2002), von Arnold et al. (2005) and Simola et al. (2012). Similar to our findings on response of warmer and drier weather, Aurela et al. (2007) and Lafleur and Humphreys (2008) also found increased GPP_{ff} with warmer growing season temperature but reduced GPP_{ff} and enhanced R_{ff} at extreme temperature in a sub-arctic fen. Our findings together with others (e.g. Griffis et al., 2000; Bubier et al., 2003; Aurela et al., 2007; Wieder et al., 2009) demonstrate the important interaction between temperature and water availability for GPP_{ff} and R_{ff} response, as either factor alone could not determine the overall growth response of peatland vegetation under changing climatic conditions. Persistently deep water table at the drained site likely limited any response to the short term drying in 2012 as this did little to further lower the water table.

Ten years of drainage in a dry continental boreal bog had a significant impact on the plant community, plant biomass and carbon fluxes, and the responses of the peatland to drainage varied between microforms and over time. Drainage replaced mosses with shrubs at hummocks and lichens at hollows such that the ground-layer aboveground biomass increased

significantly (Fig. 3). The aboveground biomass appears to be within a range of previous reports for similar types of peatlands. Published data for aboveground tree biomass across 20 bogs and ground-layer shrubs biomass across 16 bogs varied quite broadly with means of 2177 g m^{-2} ($\pm 2259 \text{ g m}^{-2}$) and 478 g m^{-2} ($\pm 224 \text{ g m}^{-2}$), respectively (Moore et al., 2002). Our data for average total of the ground-layer and aboveground tree biomass ($3490 \pm 263 \text{ g m}^{-2}$) fall within the range of the published values. The drainage-induced increase in ground-layer biomass including above- and belowground biomass observed was also reported by Moore et al. (2002). We could measure belowground biomass to only 20 cm depth due to frozen lower layers of soil at the time of sampling and therefore it is likely that we may have underestimated the root biomass particularly at the drained site with large oxic zone. However, this still likely captured the majority of belowground biomass as Lieffers and Rothwell (1987) found only 6% of root biomass occurred below 20 cm deep in a drained bog.

Although aboveground tree biomass was slightly higher at control site due to denser but smaller diameter trees, we found higher total biomass at the drained site due to its significantly higher ground-layer biomass than that at the control site. In both of the study years, the tree productivity was significantly higher at the drained site than that at control. The higher belowground biomass supported with higher R_f at the drained site, is a strong indication that lowered water table enhanced tree growth as concluded by (Hanson et al., 2000), Hermle et al. (2010) and Lieffers and Rothwell (1987). Although we determined the contribution of R_f to R_{ff} in our treed peatland study, our main aim was to quantify and include R_{ff} in seasonal model construction. The drainage induced significantly higher coverage of vascular plants and ground-layer aboveground biomass offsets some of the loss of CO_2 due to deeper oxic zone and higher decomposition rates as the water table drops (Ise et al., 2008). However, our carbon balance estimates suggest that drainage has led to a shift from CO_2 sink to a CO_2 source as the drainage-induced increase in R_{ff} (supported by R_f) was substantially higher than that of increase in GPP in both study seasons of 2011 and 2012. Similarly, Chivers et al. (2009) conducted a water table drawdown response experiment in an Alaskan moderately rich, treeless fen and found after two years of drainage, similar to our finding, that the drainage shifted the peatland from a sink of CO_2 to a source, although this change was much smaller than that of the change observed in our study of effects of drainage after 10 yr.

Peatland microforms have been shown to have different rates of CO_2 exchange and respond differently to changes in environmental conditions. For example, Waddington and Roulet (2000) found significantly higher uptake of CO_2 at a wetter microform (lawn) than that at the drier one (ridge) in over two growing seasons. Strack et al. (2006) studied CO_2 exchange following water table drawdown along a microtopographic gradient in a cool temperate poor fen and com-

pared results to a natural microtopographic gradient over two growing seasons. They also reported higher uptake of CO_2 at the wetter microform (hollow). They found that drained hummocks had lower GPP_{max} than drained hollows in contrast to control microforms and suggested that lower water tables would result in flattening of the peatland microtopography (i.e. hummocks shrink while hollows accumulate peat). In contrast, in the present study in a dry continental boreal treed bog, we found that after a decade of drainage the GPP_{max} was in fact the highest at drained hummocks in both growing seasons. The increase in GPP_{max} at drained hummocks was probably due to enhanced growth and greater coverage of shrubs. Conversely, replacement of *Sphagnum* by lichens at drained hollows (over 30 times higher biomass than at control hollows) probably led to the observed reduction in GPP_{max} (Table 4). Moreover, the drained hollows were the largest source of CO_2 in both years. Therefore we expect an increase in relative equilibrium peat depth at the hummocks and decrease in equilibrium peat depth at the hollows as an effect of drainage over the long run. These findings are not consistent with Strack et al. (2006) and are likely due to contrasting climate conditions of the two studies. For example, the earlier study was conducted in an open poor fen where average growing season precipitation recorded during the two study years were 433 and 358 mm in contrast to 402 and 281 mm recorded at our treed continental bog. Also the water table in the earlier study was much shallower and linked to regional hydrology whereas the much deeper water table in this study was controlled by the precipitation and the local elevation. On the other hand, these results together are consistent with a general “humpbacked” relationship between peat accumulation and water table depth (e.g., Belyea and Clymo, 2001; Belyea, 2009). Given the initially dry conditions at this continental bog, further drying is expected to shift both hummocks and hollows to lower rates of peat accumulation whereas a flattening of the curve at deep water tables (reduced effect of water table change on peat accumulation at drier sites) would reduce this effect at already dry sites (e.g. hummocks).

To sum up, the drained continental bog compared with a natural one simulated the potential climate-induced lowered water table and revealed spatial and temporal heterogeneity in CO_2 fluxes and plant biomass in the treed peatland complex. Drainage affected vegetation coverage, plant biomass and CO_2 fluxes differently at the microforms after a decade. Significant replacement of mosses with shrubs at hummocks and lichens at hollows increased ground-layer aboveground biomass significantly at the hummocks and generally at the hollows. This drainage-induced change in vegetation coverage and biomass shifted the bog from a sink of CO_2 to a source despite an increase in tree productivity. Net emission of CO_2 can decelerate the rate of vertical growth of microforms, whereas net uptake of CO_2 can accelerate the rate of vertical growth (Belyea, 2009). In this study we noticed significant increase in net CO_2 uptake at hummocks and net

release at hollows as a result of 10 yr of drainage (Table 4) in contrast to previous studies in wetter climates. This illustrates the importance of initial climatic conditions for predicting peatland response (e.g. Hilbert et al., 2000). Continued low water tables could lead to further shifts in vegetation in the future and thus a different CO₂-C balance than determined following 10 yr of water table drawdown.

5 Conclusions

Ten years of drainage in an ombrotrophic treed bog induced ecological succession: mosses were replaced by shrubs at hummocks and lichens at hollows. The overall greater coverage of vascular plants and higher total biomass at the drained site increased the uptake of CO₂ but the loss via respiration was even higher due to peat oxidation and increased contribution of tree root respiration. The research strongly suggests that the deepening of the unsaturated zone affected C sequestration rates differently at hummocks and hollows, potentially resulting in steepened microtopographic gradient over time. Overall, drainage promoted CO₂ emissions but offset a portion of these losses by increasing total biomass in a dry continental boreal treed bog.

Acknowledgements. This research was funded by a New Faculty Award from Alberta Innovates Technology Futures to MS Sungro Horticulture provided site access. We thank Jordanna Branham, Magnus Keith, Sharif Mahmood for assistance with fieldwork, Tak Fung for statistical support and Rick Smith for technical support for the meteorological station. Comments by two anonymous reviewers greatly improved the manuscript.

Edited by: T. Laurila

References

- Adkinson, A. C., Syed, K. H., and Flanagan, L. B.: Contrasting responses of growing season ecosystem CO₂ exchange to variation in temperature and water table depth in two peatlands in northern Alberta, Canada, *J. Geophys. Res.-Biogeosci.*, 116, G01004, doi:10.1029/2010jg001512, 2011.
- Alm, J., Schulman, L., Walden, J., Nykänen, H., Martikainen, P. J., and Silvola, J.: Carbon balance of a boreal bog during a year with an exceptionally dry summer, *Ecology*, 80, 161–174, 1999.
- Arneth, A., Kurbatova, J., Kolle, O., Shibistova, O. B., Lloyd, J., Vygodskaya, N. N., and Schulze, E. D.: Comparative ecosystem-atmosphere exchange of energy and mass in a European Russian and a central Siberian bog II. Interseasonal and interannual variability of CO₂ fluxes, *Tellus B*, 54, 514–530, 2002.
- Aurela, M., Riutta, T., Laurila, T., Tuovinen, J. P., Vesala, T., Tuittila, E. S., Rinne, J., Haapanala, S., and Laine, J.: CO₂ exchange of a sedge fen in southern Finland—the impact of a drought period, *Tellus B*, 59, 826–837, 2007.
- Belyea, L. R.: Nonlinear dynamics of peatlands and potential feedbacks on the climate system, in: *Carbon Cycling in Northern Peatlands*, Geophys. Monogr. Ser., AGU, Washington, DC, 5–18, 2009.
- Belyea, L. R. and Clymo, R.: Feedback control of the rate of peat formation, *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268, 1315–1321, 2001.
- Blodau, C. and Siems, M.: Drainage-induced forest growth alters belowground carbon biogeochemistry in the Mer Bleue bog, Canada, *Biogeochemistry*, 107, 107–123, 2012.
- Boone, R. D., Nadelhoffer, K. J., Canary, J. D., and Kaye, J. P.: Roots exert a strong influence on the temperature sensitivity of soil respiration, *Nature*, 396, 570–572, 1998.
- Bowden, R. D., Nadelhoffer, K. J., Boone, R. D., Melillo, J. M., and Garrison, J. B.: Contributions of above ground litter, below ground litter, and root respiration to total soil respiration in a temperate mixed hardwood forest, *Can. J. Forest Res*, 23, 1402–1407, 1993.
- Brant, J. B., Myrold, D. D., and Sulzman, E. W.: Root controls on soil microbial community structure in forest soils, *Oecologia*, 148, 650–659, 2006.
- Bubier, J. L., Bhatia, G., Moore, T. R., Roulet, N. T., and Lafleur, P. M.: Spatial and temporal variability in growing-season net ecosystem carbon dioxide exchange at a large peatland in Ontario, Canada, *Ecosystems*, 6, 353–367, 2003.
- Cai, T., Flanagan, L. B., and Syed, K. H.: Warmer and drier conditions stimulate respiration more than photosynthesis in a boreal peatland ecosystem: Analysis of automatic chambers and eddy covariance measurements, *Plant Cell and Environment*, 33, 394–407, 2010.
- Camill, P.: Patterns of boreal permafrost peatland vegetation across environmental gradients sensitive to climate warming, *Can. J. Botany*, 77, 721–733, 1999.
- Carlson, M., Chen, J., Elgie, S., Henschel, C., Montenegro, A., Roulet, N. T., Scott, N., Tarnocai, C., and Wells, J.: Maintaining the role of Canada's forests and peatlands in climate regulation, *The Forestry Chronicle*, 86, 434–443, 2010.
- Chapman, S. J., and Thurlow, M.: Peat respiration at low temperatures, *Soil Biol. Biochem.*, 30, 1013–1021, 1998.
- Charman, D. and Chichester: *Peatlands and Environmental Change*, John Wiley and Sons Ltd, Chichester, 2002.
- Chivers, M., Turetsky, M., Waddington, J., Harden, J., and McGuire, A.: Effects of experimental water table and temperature manipulations on ecosystem carbon dioxide fluxes in an Alaskan rich fen, *Ecosystems*, 12, 1329–1342, 2009.
- Clymo, R. S.: The growth of Sphagnum: methods of measurement, *J. Ecol.*, 58, 13–49, 1970.
- Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, 2006.
- Díaz Pinés, E., Schindlbacher, A., Pfeffer, M., Jandl, R., Zechmeister-Boltenstern, S., and Agustín, R.: Root trenching: a useful tool to estimate autotrophic soil respiration? A case study in an Austrian mountain forest, *Europ. J. Forest Res.*, 125, 101–109, 2010.
- Edwards, N. T. and Norby, R. J.: Below-ground respiratory responses of sugar maple and red maple samplings to atmospheric CO₂ enrichment and elevated air temperature, *Plant Soil*, 206, 85–97, 1999.

- Gorham, E.: Northern Peatlands: Role in the carbon cycle and probable responses to climatic warming, *Ecol. Appl.*, 1, 182–195, 1991.
- Griffis, T. J., Rouse, W. R., and Waddington, J. M.: Interannual variability of net ecosystem CO₂ exchange at a subarctic fen, *Global Biogeochemical Cycles*, 14, 1109–1121, 2000.
- Grigal, D. F. and Kernik, L. K.: Generality of black spruce biomass estimation equations, *Can. J. Forest Res.*, 14, 468–470, 1984.
- Gruber, N., Friedlingstein, P., B, F. C., Valentini, R., Heimann, M., Richey, J. E. P., Romero-Lankao, Schulze, E.-D., and Chen, C.-T. A.: The vulnerability of the carbon cycle in the 21st Century: An assessment of carbon-climate-human interactions, *The Global Carbon Cycle: Integrating Humans, Climate, and the Natural World*, Washington, DC, 2004.
- Hanson, P. J., Edwards, N. T., Garten, C. T., and Andrews, J. A.: Separating root and soil microbial contributions to soil respiration: A review of methods and observations, *Biogeochemistry*, 48, 115–146, 2000.
- Heikurainen, L. and Pakarinen, P.: Peatland classification. In *Peatlands and their utilization in Finland*, Finnish National Committee of the International Peat Society, Helsinki, 1982.
- Hengeveld, H. G.: Projections for Canada's climate future, *Climate change digest*, 500, 1–27, 2000.
- Hermle, S., Lavigne, M. B., Bernier, P. Y., Bergeron, O., and Pare, D.: Component respiration, ecosystem respiration and net primary production of a mature black spruce forest in northern Quebec, *Tree Physiology*, 30, 527–540, 2010.
- Hilbert, D. W., Roulet, N., and Moore, T.: Modelling and analysis of peatlands as dynamical systems, *Journal of Ecology*, 88, 230–242, 2000.
- IPCC: Impacts, Adaptation and Vulnerability: Contribution of working group II to the fourth assessment report of the inter-governmental panel on climate change, edited by: Parry, M. L., Canziani, O. F., Palutikof, J. P., van der Linden, P. J., and Hanson, C. E., Cambridge University Press, Cambridge, United Kingdom, 976 pp., 2007.
- Ise, T., Dunn, A. L., Wofsy, S. C., and Moorcroft, P. R.: High sensitivity of peat decomposition to climate change through water-table feedback, *Nat. Geosci.*, 1, 763–766, 2008.
- Janssens, I. A., Lankreijer, H., and Matteucci, G.: Productivity overshadows temperature in determining soil and ecosystem respiration across European forests, *Glob. Change Biol.*, 269–278, 2001.
- Joosten, H. and Clarke, D.: Wise use of mires and peatlands, *International mire conservation group*, 2002.
- Koskinen, M., Minkinen, K., Ojanen, P., Kämäräinen, M., Laurila, T., and Lohila, A.: Measurements of CO₂ exchange with an automated chamber system throughout the year: challenges in measuring nighttime respiration on porous peat soil, *Biogeosciences Discuss.*, 10, 14195–14238, 2013, <http://www.biogeosciences-discuss.net/10/14195/2013/>.
- Kutzbach, L., Schneider, J., Sachs, T., Giebels, M., Nykänen, H., Shurpali, N. J., Martikainen, P. J., Alm, J., and Wilkening, M.: CO₂ flux determination by closed-chamber methods can be seriously biased by inappropriate application of linear regression, *Biogeosciences*, 4, 1005–1025, doi:10.5194/bg-4-1005-2007, 2007.
- Kuz'yakov, Y.: Sources of CO₂ efflux from soil and review of partitioning methods, *Soil Biol. Biochem.*, 425–448, 2006.
- Lafleur, P. M. and Humphreys, E. R.: Spring warming and carbon dioxide exchange over low Arctic tundra in central Canada, *Glob. Change Biol.*, 14, 740–756, 2008.
- Lai, D. Y. F., Roulet, N. T., Humphreys, E. R., Moore, T. R., and Dalva, M.: The effect of atmospheric turbulence and chamber deployment period on autochamber CO₂ and CH₄ flux measurements in an ombrotrophic peatland, *Biogeosciences*, 9, 3305–3322, doi:10.5194/bg-9-3305-2012, 2012.
- Li, Z., Kurz, W. A., Apps, M. J., and Beukema, S. J.: Belowground biomass dynamics in the Carbon Budget Model of the Canadian Forest Sector: recent improvements and implications for the estimation of NPP and NEP, *Can. J. Forest Res.*, 33, 126–136, 2003.
- Lieffers, V. and Rothwell, R.: Rooting of peatland black spruce and tamarack in relation to depth of water table, *Can. J. Botany*, 65, 817–821, 1987.
- Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H., and Schaeppman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications – a synthesis, *Biogeosciences*, 5, 1475–1491, doi:10.5194/bg-5-1475-2008, 2008.
- Lohila, A., Minkinen, K., Aurela, M., Tuovinen, J.-P., Penttilä, T., Ojanen, P., and Laurila, T.: Greenhouse gas flux measurements in a forestry-drained peatland indicate a large carbon sink, *Biogeosciences*, 8, 3203–3218, doi:10.5194/bg-8-3203-2011, 2011.
- Loisel, J., Gallego-Sala, A. V., and Yu, Z.: Global-scale pattern of peatland Sphagnum growth driven by photosynthetically active radiation and growing season length, *Biogeosciences*, 9, 2737–2746, doi:10.5194/bg-9-2737-2012, 2012.
- Macdonald, S. E. and Lieffers, V. J.: Photosynthesis, water relations, and foliar nitrogen of *Picea mariana* and *Larix laricina* from drained and undrained peatlands, *Can. J. Forest Res.*, 20, 995–1000, 1990.
- Moore, P. D.: The future of cool temperate bogs, *Environmental Conservation*, 29, 3–20, 2002.
- Moore, T. R., and Dalva, M.: The influence of temperature and water table position on carbon dioxide and methane emissions from laboratory columns of peatland soils, *J. Soil Sci.*, 44, 651–664, 1993.
- Moore, T. R., Bubier, J. L., Frohling, S. E., Lafleur, P. M., and Roulet, N. T.: Plant biomass and production and CO₂ exchange in an ombrotrophic bog, *J. Ecol.*, 90, 25–36, 2002.
- Moore, T. R., Lafleur, P. M., Poon, D. M. I., Heumann, B. W., Seaquist, J. W., and Roulet, N. T.: Spring photosynthesis in a cool temperate bog, *Global Change Biol.*, 12, 2323–2335, 2006.
- Mullin, T., Morgenstern, E., Park, Y., and Fowler, D.: Genetic parameters from a clonally replicated test of black spruce (*Picea mariana*), *Can. J. Forest Res.*, 22, 24–36, 1992.
- Ow, L. F., Griffin, K. L., Whitehead, D., Walcroft, A. S., and Turnbull, M. H.: Thermal acclimation of leaf respiration but not photosynthesis in *Populus deltoides* × *nigra*, *New Phytologist*, 178, 123–134, 2008.
- Parmentier, F., Van der Molen, M., de Jeu, R., Hendriks, D., and Dolman, A.: CO₂ fluxes and evaporation on a peatland in the Netherlands appear not affected by water table fluctuations, *Agr. Forest Meteorol.*, 149, 1201–1208, 2009.
- Price, J. S.: The Hydrology of Peatlands of Eastern Canada, in: *Ecologie des tourbières du Québec et Labrador*, edited by: Payette, S. a. R., L, 141–158, 2010.

- Rey, A., Pegoraro, E., Tedeschi, V., De-Parri, I., Jarvis, P. G., and Valentini, R.: Annual variation in soil respiration and its components in a coppice oak forest in Central Italy, *Glob. Change Biol.*, 851–866, 2002.
- Riley, J. L.: *The Vascular Flora of the Hudson Bay Lowland and Its Postglacial Origins*, NRC Research Press, 2003.
- Riutta, T., Laine, J., Aurela, M., Rinne, J., Vesala, T., Laurila, T., Haapanala, S., Pihlatie, M., and Tuittila, E. S.: Spatial variation in plant community functions regulates carbon gas dynamics in a boreal fen ecosystem, *Tellus B*, 59, 838–852, 2007.
- Robroek, B. J. M., Schouten, M. G. C., Limpens, J., Berendse, F., and Poorter, H.: Interactive effects of water table and precipitation on net CO₂ assimilation of three co-occurring Sphagnum mosses differing in distribution above the water table, *Glob. Change Biol.*, 15, 680–691, 2009.
- Roulet, N.: Peatlands, carbon storage, greenhouse gases, and the Kyoto Protocol: Prospects and significance for Canada, *Wetlands*, 20, 605–615, 2000.
- Roulet, N. T., Moore, T. I. M., Bubier, J., and Lafleur, P.: Northern fens: methane flux and climatic change, *Tellus B*, 44, 100–105, 1992.
- Roulet, N. T., Lafleur, P. M., Richard, P. J. H., Moore, T. R., HumphreySk, E. R., and Bubier, J.: Contemporary carbon balance and late Holocene carbon accumulation in a northern peatland, *Glob. Change Biol.*, 13, 397–411, 2007.
- Ryan, M. G.: Foliar maintenance respiration of subalpine and boreal trees and shrubs in relation to nitrogen content, *Plant, Cell and Environ.*, 18, 765–772, 1995.
- Simola, H., Pitkänen, A., and Turunen, J.: Carbon loss in drained forestry peatlands in Finland, estimated by re-sampling peatlands surveyed in the 1980s, *Europ. J. Soil Sci.*, 63, 798–807, 2012.
- Strack, M., Waddington, J., Rochefort, L., and Tuittila, E. S.: Response of vegetation and net ecosystem carbon dioxide exchange at different peatland microforms following water table draw-down, *J. Geophys. Res.* (2005–2012), 111, G02006, 2006.
- Syed, K. H., Flanagan, L. B., Carlson, P. J., Glenn, A. J., and Van Gaalen, K. E.: Environmental control of net ecosystem CO₂ exchange in a treed, moderately rich fen in northern Alberta, *Agr. Forest Meteorol.*, 140, 97–114, 2006.
- Szumigalski, A. R. and Bayley, S. E.: Net above-ground primary production along a bog-rich fen gradient in central Alberta, Canada, *Wetlands*, 16, 467–476, 1996.
- Tarnocai, C.: The effect of climate change on carbon in Canadian peatlands, *Glob. Planet. Change*, 53, 222–232, 2006.
- Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Glob. Biogeochem. Cy.*, 23, GB003327, doi:10.1029/2008gb003327, 2009.
- Thormann, M. N. and Bayley, S. E.: Aboveground net primary production along a bog-fen-marsh gradient in southern boreal Alberta, Canada, *Ecoscience*, 4, 374–384, 1997.
- Tokida, T., Mizoguchi, M., Miyazaki, T., Kagemoto, A., Nagata, O., and Hatano, R.: Episodic release of methane bubbles from peatland during spring thaw, *Chemosphere*, 70, 165–171, doi:10.1016/j.chemosphere.2007.06.042, 2007.
- Tuittila, E. S., Vasander, H., and Laine, J.: Sensitivity of C Sequestration in Reintroduced Sphagnum to Water-Level Variation in a Cutaway Peatland, *Rest. Ecol.*, 12, 483–493, 2004.
- Turetsky, M., Wieder, K., Halsey, L., and Vitt, D.: Current disturbance and the diminishing peatland carbon sink, *Geophys. Res. Lett.*, 10, 131 pp., doi:10.1029/2001GL014000, 2002.
- Turunen, J., Tomppo, E., Tolonen, K., and Reinikainen, A.: Estimating carbon accumulation rates of undrained mires in Finland—application to boreal and subarctic regions, *The Holocene*, 12, 69–80, 2002.
- Updegraff, K., Bridgham, S. D., Pastor, J., Weishampel, P., and Harth, C.: Response of CO₂ and CH₄ emissions from peatlands to warming and water table manipulation, *Ecol. Appl.*, 11, 311–326, 2001.
- Valentini, R., Matteucci, G., and Dolman, A. J.: Respiration as the main determinant of carbon balance in European forests, *Nature*, 861–865, 2000.
- Varem-Sanders, T. M. L. and Campbell, I. D.: DendroScan: a tree-ring width and density measurement system. Canadian Forest Service, Northern Forestry Centre, Edmonton, Alta. Spec. Rep., 10, 131 pp., 1996.
- Vitt, D. H., Halsey, L. A., Thormann, M. N., and Martin, T.: Peatland Inventory of Alberta. Phase 1: Overview of Peatland Resources in the Natural Regions and Subregions of the Province. Sustainable Forest Management Network, University of Alberta, Edmonton, Alberta, 1998.
- Vitt, D. H., Wieder, R. K., Scott, K. D., and Faller, S.: Decomposition and peat accumulation in rich fens of boreal Alberta, Canada, *Ecosystems*, 12, 360–373, 2009.
- von Arnold, K., Hånell, B., Stendahl, J., and Klemedtsson, L.: Greenhouse gas fluxes from drained organic forestland in Sweden, *Scand. J. Forest Res.*, 20, 400–411, 2005.
- Waddington, J., Warner, K., and Kennedy, G.: Cutover peatlands: a persistent source of atmospheric CO₂, *Global Biogeochem. Cy.*, 16, 1002, doi:10.1029/2001GB001398, 2002.
- Waddington, J. M. and Price, J. S.: Effect of peatland drainage, harvesting, and restoration on atmospheric water and carbon exchange *Physical Geography [Phys. Geogr.]*, 21, 433–451, 2000.
- Waddington, M. J. and Roulet, N. T.: Carbon balance of a boreal patterned peatland., *Glob. Change Biol.*, 6, 87–97, 2000.
- Wang, X., Zhu, B., Wang, Y., and Zheng, X.: Field measures of the contribution of root respiration to soil respiration in an alder and cypress mixed plantation by two methods: trenching method and root biomass regression method, *Europ. J. Forest Res.*, 127, 285–291, 2008.
- Weltzin, J. F., Pastor, J., Harth, C., Bridgham, S. D., Updegraff, K., and Chapin, C. T.: Response of bog and fen plant communities to warming and water-table manipulations, *Ecology*, 81, 3464–3478, 2000.
- Wieder, R. K., Scott, K. D., Kamminga, K., Vile, M. A., Vitt, D. H., Bone, T., Xu, B., Benscotter, B. W., and Bhatti, J. S.: Postfire carbon balance in boreal bogs of Alberta, Canada, *Glob. Change Biol.*, 15, 63–81, 2009.